



Contents lists available at ScienceDirect

Journal of Advanced Research

journal homepage: www.elsevier.com/locate/jare

Original Manuscript

Exerkine irisin mitigates cognitive impairment by suppressing gut-brain axis-mediated inflammation

Hu Zhang¹, Jiling Liang¹, Jielun Huang, Minghui Wang, Liangwen Wu, Tong Wu^{*}, Ning Chen^{*}


Tianjiu Research and Development Center for Exercise Nutrition and Foods, Hubei Key Laboratory of Exercise Training and Monitoring, College of Sports Medicine, Wuhan Sports University, Wuhan 430079, China

HIGHLIGHTS

- Irisin can improve learning and memory capacity of aged mice.
- Irisin can reverse disturbed gut microbiota in aged mice.
- Irisin can optimize metabolites of gut microbiota in aged mice.
- Exercise-induced irisin as a mediator regulating gut microbiota and metabolites for suppressing gut-brain axis-mediated inflammation.

GRAPHICAL ABSTRACT

Exercise-induced irisin ameliorates cognitive impairment of aged mice through rescuing the imbalanced gut microbiota and suppressing TLR4/MyD88 signal pathway-mediated insulin resistance.



ARTICLE INFO

Article history:

Received 3 February 2024

Revised 16 April 2024

Accepted 26 October 2024

Available online xxx

Keywords:

Exercise

Irisin

Gut microbiota

Cognitive capacity

Gut-brain axis

ABSTRACT

Introduction: Exercise has been recognized to improve cognitive performance by optimizing gut flora and up-regulating exerkine irisin.

Objective: Although exercise-induced irisin is beneficial to cognitive improvement, whether this benefit is achieved by optimizing gut microbiota and metabolites is not fully explored.

Methods: After aerobic exercise and exogenous irisin interventions for 12 weeks, the 16S rRNA and metabolites in feces of 21-month-old mice were analyzed. Meanwhile, the differential miRNAs and mRNAs in hippocampal tissues were screened by high-throughput sequencing. Relevant mRNAs and proteins were evaluated by RT-PCR, Western blot, and immunofluorescence.

Results: Compared with the young control mice, irisin levels and cognitive capacity of aged mice revealed a significant reduction, while aerobic exercise and intraperitoneal injection of exogenous irisin reversed aging-induced cognitive impairment. Similarly, 147 up-regulated and 173 down-regulated metabolites were detected in aged mice, while 64 and 45 up-regulated and 225 and 187 down-regulated metabolites were detected in aged mice with exercise and irisin interventions, respectively. Moreover, during

* Corresponding authors.

E-mail addresses: 2021668@whsu.edu.cn (T. Wu), nchen510@gmail.com (N. Chen).

¹ These authors have contributed equally to this work.

<https://doi.org/10.1016/j.jare.2024.10.031>

2090-1232/© 2024 Published by Elsevier B.V. on behalf of Cairo University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

hippocampal miRNA and mRNA sequencing analysis, 9 differential gut flora and 35 differential genes were identified to be correlated with the inflammatory signaling mediated by the TLR4/MyD88 signal pathway.

Conclusion: Aging-induced cognitive impairment is due to insulin resistance induced by TLR4/MyD88 signaling activation in hippocampal tissues mediated by gut microbiota and metabolite changes. Myokine irisin may be an important mediator in optimizing gut microbiota and metabolism for an improved understanding of mitigated aging process upon exercise interventions.

© 2024 Published by Elsevier B.V. on behalf of Cairo University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Cognitive decline, as an important manifestation of aging, is common in neurodegenerative diseases such as Alzheimer's disease (AD), with cognitive impairment [1,2]. The global population aging reveals a rapid expansion with increasing number of the elderly suffering from cognitive decline. With the in-depth studies, more and more factors including aging, increased inflammation [3], impaired autophagy [4], excessive endoplasmic reticulum (ER) stress [5], abnormal circadian rhythm [6], and imbalanced gut microbiota [7] are involved in cognitive decline. Exercise as a non-pharmacological intervention can promote body health involving complex molecular mechanisms. It is worth noting that imbalanced gut microbiota can be induced by aging process [8] and the mitigation of cognitive decline through optimizing gut microbiota and their metabolites has been verified [9], which could become one of the key interventional targets for repealing global population aging [10].

Recent studies have confirmed that exercise-induced myokine irisin can enhance cognitive and memory capacity of aged mice; however, underlying mechanisms for alleviating impaired cognition through exercise-induced irisin-mediated optimization of gut microbiota and metabolites have less reported [11]. Irisin has been found to be involved in the regulation of intestinal microbiota in myocardial ischemia and colitis protection, and an obvious depression and a significantly cognitive decline have been observed in animal models with irisin deficiency [11–13]. Similarly, another study has demonstrated that irisin may exert anti-inflammatory effects by targeting the TLR4/MyD88 signal pathway [14]. As we all know, TLR4 is mainly activated by lipopolysaccharide (LPS) in the intestine, thereby impairing the balance of gut flora in the body. Coincidentally, during the aging process, the gut flora could be imbalanced, thereby resulting in the increased intestinal epithelial permeability, leading to an increased LPS level in the body, stimulating inflammatory factors, aggravating central nervous system inflammation, and triggering insulin resistance and cognitive impairment [15–17]. In a model of type 1 diabetes, injection of exogenous irisin reveals a significant reduction of intestinal epithelial permeability [18]. Relevant studies have also confirmed that cognitive decline of aging mice is closely related to the activation of TLR4 [19], and the TLR4 knockout mouse model during aging process shows improved brain structure and cognitive capacity [20]. Therefore, it may be of great significance to target the TLR4 signal pathway for the prevention and treatment of aging-induced cognitive impairment. Similarly, we hypothesize that exercise-induced myokine irisin is likely to be involved in the regulation of intestinal microorganisms to indirectly enhance the cognitive capacity of aged mice. Through high-throughput sequencing of gut microbiota and metabolites in feces and the screening and identification of differential miRNAs and mRNAs in hippocampal tissues from aged mice followed by exercise and irisin interventions, we could deeply uncover exercise-induced irisin for rescuing aging-induced cognitive impairment as a novel interventional strategy.

Materials and methods

Recombinant irisin expression and purification

The DNA sequence encoding irisin was cloned into pET-28a vector (MilliporeSigma, CA, USA) to obtain the pET-28a-irisin plasmid. The resultant pET-28a-irisin plasmid was then transformed into *Escherichia coli* BL21(DE3) competent cells, and spread on a Luria-Bertani (LB) agar plate containing kanamycin (50 µg/L) for incubation overnight at 37 °C. A single colony was picked up to inoculate into the LB medium containing kanamycin (50 µg/L) for cell culture with 200-RPM shaking at 37 °C. Upon the optical density at 600 nm of 0.6 for cell culture, isopropyl-β-D-thiogalactopyranoside (IPTG) (0.5 mM) was added in cell culture for inducing the expression of histidine-tagged irisin during cell cultivation for another 8 h. To purify irisin, cell pellets were sonicated to disrupt cell membrane. The supernatant was filtered and injected into a nick-chelating agarose column (MilliporeSigma, CA, USA) loaded with 0.1 M nickel sulfate solution. The bound irisin protein was eluted with a gradient of imidazole from 10–500 mM in phosphate buffer. The collected protein was dialyzed in 500 mM Tris buffer (Servicebio Technology, Wuhan, China) to remove imidazole and endotoxin through dialysis membrane with the cutoff of 10000 Da. The concentration of purified recombinant irisin was determined by the BCA Protein Assay Kit (Beyotime Biotechnology, Shanghai, China) and used for the intervention of animal experiments later.

Animal grouping and interventions

Thirty 4-month-old specific pathogen-free (SPF) grade male C57BL/6 mice were fed with adequate maintenance feed and water until 21 months old, and then randomly divided the aged mice into a non-intervention group (OC), an aerobic exercise group (OE), and an irisin intraperitoneal injection group (OI), with 10 mice in each group; and ten 4-month-old mice were used as the young control group (YC). Furthermore, room temperature (25 ± 2 °C), relative humidity (50–60%), circadian rhythm (12 h:12 h), and free accessibility of foods and water were kept at same for all mice. All mice in the OE group were subjected to running training on treadmill with a slope of 10° for 12 weeks. At the beginning of running training, the mice were provided with adaptive treadmill running for 5 days (start with 4.2 m/min until 12 m/min, with an increasing interval of 1 m/min every 30 s), with exercise training time of 10, 20, 30, 45 and 45 min in first five days, respectively. After adaptive running training, the formal aerobic exercise training (12 m/min) with the training duration of 45 min and 5 times of training within one week was conducted for 12 consecutive weeks. In the OI group, the mice were provided with the intraperitoneal injection of exogenous recombinant irisin according to their body weights (500 µg/kg). Both aerobic exercise training and irisin injection were performed 5 times a week at the fixed time and the health status of all aging mice during the interventions was also recorded.

Ethics statement

All experiments involving animals were conducted according to the ethical policies and procedures approved by Institutional Animal Care and Use Committee of Wuhan Sports University with the certificate number of S087-21-05D, and complied with the internationally recognized 3R principles.

Behavioral testing

The learning and memory capacity and behavior of all mice were evaluated by Morris water maze (MWM) testing. On the 1st day, a positioning test was conducted, and the mice were released to four different locations in the east, west, south, and north directions, and allowed to freely explore and swim for 60 s to adapt to the environment. In the next 4 days, the mice were released for 4 times within one day at different starting points. If the mice did not find the platform within 60 s, they were guided to the platform and stayed on the platform for 10 s for spatial location memory. On the 5th day, after the platform was removed, the mice were released in the quadrant opposite the platform to freely swim for 60 s, and the time of targeting the platform quadrant and the number of crossing the platform of the mice were used as the indicators to evaluate the spatial learning and memory capacity.

Blood sample collection and ELISA

Prior to sample collection, all mice were fasted with free accessibility of water for 12 h, and sacrificed by carbon dioxide inhalation for collecting blood samples in sterile EP tubes from eyeballs, and the collected blood samples were stood quietly at room temperature overnight. On the next day, the blood samples were subjected to 20-min centrifugation at 3000×g and 4 °C, and the supernatant was harvested for corresponding analysis. Irisin in serum of the mice was determined by its ELISA kit (Phoenix Pharmaceuticals, California, USA), and LPS, IL-1 β , IL-6, TNF- α and LBP in serum were evaluated by corresponding ELISA kits (Meimian Industrial Co., Ltd, Jiangsu, China) in accordance with the manufacturer's operation procedures.

Collection and analysis of hippocampal tissue

Transmission electron microscope (TEM) inspection

After completing MWM testing, the mouse was sacrificed by carbon dioxide inhalation, and the hippocampal tissue at 1 mm³ was harvested, immediately fixed in 250 μ L of 2.5% glutaraldehyde containing 0.1 M sodium cacodylate (pH 7.2), subsequently post-fixed in 1% osmium tetroxide for 1 h at 4 °C, stained with uranyl acetate, and then embedded with resin. The processed block of hippocampal tissue was sectioned into 70 nm ultra-thin sections. The ultrastructure of hippocampal tissues was examined and photographed under a TEM (HT7700 Hitachi, Tokyo, Japan) at Medical and Structural Biology Research Center of Wuhan University.

Histological analysis of hippocampal tissue

Similarly, after MWM test was completed, 3 mice from each group were randomly selected on the next day, and then subjected to the perfusion using saline and 4% paraformaldehyde (pH 7.4) in the anesthesia status. The perfused mice were sacrificed by carbon dioxide inhalation, the whole brains were harvested to fix in 4% paraformaldehyde overnight, sequentially embedded in paraffin, and cut into the slices in the thickness of 5 μ m using a cryotome. After the paraffin removal, the tissue slices were subjected to hematoxylin-eosin (HE), Nissl, and immunohistochemical staining, respectively. The histological images of hippocampal tissues were acquired by imaging system (Eclipse-E100, Nikon, Japan) and analyzed by ImageJ Pro software (NIH, Bethesda, MD, USA).

Sequencing of miRNAs and mRNAs in hippocampal tissue

According to the operation procedures of the Trizol extraction kit (Sangon Biotech, Shanghai, China) and Qubit2.0 RNA detection kit (ThermoFisher, MA, USA), total RNA in hippocampal tissue was extracted and quantitatively determined. The integrity and contamination of RNA were evaluated by agarose gel.

miRNA: Total miRNA was extracted and processed for sequencing by Illumina NovaSeq6000 platform (Illumina, TX, USA). FastQC (v0.11.2) was utilized to evaluate the quality of sequencing data for all samples, the Adapter was removed and the quality was controlled by Trimmomatic. Bowtie2 (v2.2.4) software was applied for mapping the clean data to the reference genome. The miRDeep2 (v2.0.0.8) software and miRbase (v22.1) database were used to match and identify miRNAs.

mRNA: Total mRNA was extracted and processed for sequencing by Illumina NovaSeq6000 platform (Illumina, TX, USA). FastQC (v0.11.2) was used for evaluating the quality of sequencing data for all samples, the Adapter was removed and the quality was controlled by Trimmomatic (v0.36). Qualimap (v2.2.1) was used to check the distribution of uniformity and genome structure according to comparison results for further analysis. StringTie (v1.3.3b) was applied to construct known gene models for evaluating gene expression levels. Co-expression analysis of genes was conducted by WGCNA (v1.51), and multi-directional statistical analysis was performed based on sample expression matrix for sample comparison analysis. The R package DESeq (v1.26.0) was used for the analysis of the difference in gene expression, and the graph visualization was completed based on ggplot2 (v2.2.1). The interaction network between miRNAs and genes was constructed by using the igraph R package (v1.0.1) association analysis.

Dimensionality reduction analysis was implemented based on principal component analysis (PCA), the volcano plot was established to visualize differential expression, and the enrichment bubble map was generated to visualize the KEGG pathway enrichment analysis. The data in this study are mainly analyzed with the help of NCBI NR, NCBI NT, KEGG and other databases.

The analysis of 16S rRNA sequencing and gut microbiota in feces

After experimental interventions, fresh feces of the mice from all groups were collected for extracting DNA by magnetic soil and stool DNA kit (Sangon Biotech, Shanghai, China). The purity of the extracted DNA was validated by agarose gel electrophoresis. The 16S rRNA gene was amplified by PCR and purified by 2% agarose gel electrophoresis. The target band was recovered by the Universal DNA Purification and Recovery Kit (ThermoFisher, MA, USA). NEB Next® Ultra DNA Library Prep Kit (Bio-Rad Laboratories, San Francisco, USA) was used for the construction of gene library. After the library was qualified, the V3-V4 region of 16S rRNA was sequenced using the NovaSeq6000 sequencing platform (Illumina, TX, USA). Quality of raw data was evaluated by FastQC (v0.11.2), Trimmomatic (v0.36) was used for removing adapter and quality control, and Flash (v1.2.7) was used for splice data to obtain clean data. Qiime (v1.9.1) was used to denoise, cluster, and annotate with species. Then, α -diversity and β -diversity analysis (Permanova test), cluster analysis (PCA analysis), difference analysis (LEfSe), and correlation analysis of the species were carried out at each taxonomic level.

Sequencing of non-targeted metabolites in gut

Fecal samples were thawed on ice after being removed from the -80 °C freezer, and approximately 20 mg of the samples were weighed for metabolite extraction. Non-targeted metabolites were collected using ultra-high performance liquid chromatography

(UPLC) (SCIEX, MA, USA) and quadrupole-time-of-flight mass spectrometry (Q-TOF-MS) (TripleTOF6600, AB SCIEX, MA, USA) for data acquisition. Extensive targeted detection data were acquired by UPLC (SCIEX, MA, USA) and tandem mass spectrometry (MS/MS) (SCIEX, MA, USA). The data was processed by Progenesis QI software (v3.0), and the data was imported into EZinfo3.0 (v3.0.3) software for multidimensional statistical analysis. Metabolites were analyzed by PCA (R-Base Package v3.5.1) and orthogonal partial least squares discriminant analysis (OPLS-DA) (MetaboAnalystR v1.0.1) for multidimensional statistical analysis between two groups. The differential metabolites with vasoactive intestinal peptide (VIP) > 1 were screened and subjected to unidimensional statistical analysis (Welch's *t*-test and fold change analysis). Moreover, differential metabolites meeting the criteria (VIP > 1, $p < 0.05$ and fold change > 2) were screened and subjected to final compound identification analysis. The HMDB and METLIN databases were searched based on the primary and secondary spectra to obtain the identification results of differential metabolites. MetaboAnalyst (v5.0) online software was used to analyze the signal pathway and enrichment of differential metabolites.

qRT-PCR

RNA-easy isolation reagent was used for extracting mRNA in hippocampal tissues. HiSipt® II Q RT Supermix (Vazyme, Nanjing, China) was used for qRT-PCR to reverse-transcribe mRNA into cDNA after removing genomic DNA. Taq Pro Universal SYBR qPCR Master Mix (Vazyme, Nanjing, China) was used for qRT-PCR of mRNA. The primers were synthesized from Sangon Biotech (Table 1). Quant Studio 1 Real-Time PCR System (ThermoFisher, MA, USA) was used to perform qRT-PCR with 20 μ L reaction system for each sample in triple repeats using β -actin as the control, and the $2^{-\Delta\Delta CT}$ method was performed for all calculations.

Western blot

Hippocampal tissue samples from the mice were collected and immediately frozen in liquid nitrogen, and then stored in -80°C refrigerator for future evaluation. During western blot analysis, frozen hippocampal tissue samples were taken out and homogenized in lysis buffer supplemented with protease and phosphatase inhibitors (Servicebio Biotechnology, Wuhan, China), and then extensively lysed for 30 min on ice. The lysed samples were centrifuged at $10,000\times g$ for 5 min at 4°C to obtain supernatants and the concentrations of total protein in supernatants were measured by a BCA kit (Beyotime Biotechnology, Shanghai, China). An aliquot of the supernatant was mixed with reduced protein loading buffer and heated at 95°C water bath for 5 min to denature proteins. Approximately 25 μ g of total protein in prepared samples were separated by using 8–12% sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE) and then transferred to polyvinylidene difluoride (PVDF) membrane (Merck Millipore,

MA, USA). The target protein was probed by primary antibodies against JNK, p-JNK, IL-1 β , IL-6, TNF- α , TLR4, MyD88, MKK7, p-MKK7, p38 MAPK, p-p38 MAPK, IRS, and p-IRS, as well as corresponding secondary antibodies (Cell Signaling Technology, Danvers, MA, USA). The band of the target protein after incubated with enhanced chemiluminescence (ECL) (Bio-Rad Laboratories, San Francisco, USA) reagent was imaged and visualized by an ultra-sensitive fluorescence/chemiluminescence imaging system, ChemiScope6300 (CLiNX Science Instruments, Shanghai, China).

Statistical analysis

All above experiments were accomplished according to the designated procedures (Fig. 1) and all collected data were presented as mean $\pm$ standard deviation ($M \pm SD$) to conduct the statistical analysis by GraphPad Prism software (v4.0.2). Two-tailed unpaired student's *t*-test and the Wilcoxon sum test were utilized to evaluate the statistical significance between two or multiple groups, respectively. Statistical analysis among multiple groups was implemented based on Kruskal-Wallis test for non-parametric data, and the significant difference was considered at $p < 0.05$. Spearman correlation analysis was used for correlation analysis.

Results

Exercise and irisin interventions reversed aging-induced reduction of cognitive capacity in mice

The aging-induced reduction of cognitive capacity can be reversed by aerobic exercise [21], but whether it is correlated with exercise-induced myokine irisin is still unknown. Therefore, the 21-month-old mice were provided with aerobic exercise training and intraperitoneal injection of exogenous irisin for 12 weeks, and the irisin level in serum and cognitive capacity of the mice were evaluated. The irisin level in the OC group were decreased significantly ($p = 0.0154$) when compared with the YC group. Oppositely, compared with the OC group, the irisin in OE and OI groups exhibited an obvious increase ($p = 0.0134$, $p = 0.0037$). Notably, the irisin level in the OI group was higher than that in the OE group (Fig. 2a). Based on cognition evaluation of the mice through MWM testing, compared with the YC group, the escape latency to platform increased significantly ($p = 0.0213$), while the target quadrant-exploring time and the platform-crossing number of the mice from the OC group on the 5th day were markedly reduced ($p = 0.0067$, $p = 0.0021$); in contrast, the reduced escape latency to platform, and the increased target quadrant-exploring time and platform-crossing number of the mice from OE and OI groups on the 5th day were observed ($p = 0.0041$, $p = 0.0425$; $p = 0.0005$, $p = 0.0491$; $p = 0.0002$, $p = 0.0075$) (Fig. 2c-e). These results demonstrated that 24-month-old mice exhibited the significant down-regulation of irisin in serum and the reduced cognitive capacity, which could be reversed by aerobic exercise and intraperitoneal injection of exogenous irisin.

Exercise and irisin interventions rescued aging-induced damage of hippocampal tissues in mice

Aging process is accompanied by damaged morphology, dysfunction, and reduced neuron number of hippocampal tissues; in contrast, exercise can effectively reverse these changes, as confirmed by HE and Nissl staining and immunohistochemical analysis. Hippocampal tissues of the mice from the OC group had sparse neurons and unclear nucleoli and nuclei, as shown in darker image by Nissl staining (Fig. 3a and b), and significantly reduced the number of neurons in CA1, CA3 and DG areas when compared with the YC group ($p = 0.0015$, $p = 0.0076$, $p = 0.0009$) (Fig. 3c). On the

Table 1
Primer information for qRT-PCR.

Gene name	Forward primer	Reverse primer
LBP	GGCTTGGCGTGGTCACTAA	TGCCGACTTTGGATTTCGATCA
TLR4	AGATCTGAGCTTCAACCCCT	TGTTTCAATTTACACCTGGATAA
MyD88	CTCGCAGTTTGTGGATGCC	TTTCTGGCAGTCTCTCTCGAT
MKK7	CGCGTCTCTGGTTAAGGATGT	CTCCAGACTCCCACTGAAGAA
JNK	GAACAGGATTGAGTAGCGGC	ATATTACCAAGCGCGCAG
p38 MAPK	AAAGGACCTACCGAGAGTTGC	GTCACCAGGTACACGTCTATT
NF- κ B	CCGAATCTCTGGACAGCTT	GTTTCGAGTAGCCATACCTGG
TNF- α	CCAAATGGCTCCCTCTCAT	CCACTTGGTGGTTTGCTACGA
IL-1 β	GCTTCCTTGTGCAAGTGTCT	TGACCACTCTCCAGTACCCA
IL-6	GACTGGGGATGTCTGTAGCTC	CAACTGGATGGAAGTCTCTTGC




Fig. 1. Schematic diagram of the experimental process.




Fig. 2. Irisin levels in serum and cognitive capacity of the mice from different groups. (a) Irisin levels in serum of the mice from different groups; (b) Swimming track during MWM testing; (c) The escape latency to platform during MWM testing; (d) The proportion of target quadrant-exploring time of the mice on the fifth day; (e) The number of crossing platform of the mice on the fifth day.




Fig. 3. HE and Nissl staining, and TEM observation of neurons in hippocampal tissues of the mice from each group. (a) HE staining and (b) Nissl staining of hippocampal tissues in the mice from different groups; (c) Immunohistochemical staining of mature neurons; (d) The number of mature neurons in different regions of hippocampal tissues; (e) TEM observation of hippocampal tissues; (f) The number of synapses; (g) The length of synaptic arcs; (h) The thickness of postsynaptic density; (i) The width of protrusion gaps. Red arrows indicate synapses.

contrary, compared with the OC group, the neurons in CA1, CA3 and DG areas of hippocampal tissues in mice from OE and OI groups revealed more ordered and dense arrangement, and less damage and more matured neurons ($p = 0.0363$, $p = 0.0163$, $p = 0.0008$; $p = 0.0018$, $p = 0.0058$) (Fig. 3d). These results suggested that neurons in CA1, CA3 and DG areas in hippocampal tissues of aged mice were disordered and sparse during aging process, and the aging-induced reduction of neuronal function and number could be reversed by aerobic exercise and intraperitoneal injection of exogenous irisin.

Similarly, aging accelerated the loss of neuronal synapses, increased protrusion gaps, reduced synaptic arc lengths, and decreased postsynaptic density, the hallmarks of synaptic plasticity, as shown in TEM images (Fig. 3e). Indeed, based on the statistical analysis, the number of neuronal synapses decreased significantly ($p = 0.0002$) (Fig. 3f), the length of synaptic arc and the thickness of postsynaptic density decreased ($p = 0.0004$, $p = 0.0313$) (Fig. 3g and h), and synaptic gaps increased in the OC group ($p = 0.0311$) when compared with the YC group (Fig. 3i).

On the other hand, compared with the OC group, hippocampal tissues of the mice from OE and OI groups revealed significantly increased number of neurons, extended length of synaptic arcs and thickness of postsynaptic density, and reduced synaptic cleft ($p = 0.0001$, $p = 0.0012$, $p = 0.0006$, $p = 0.0003$; $p = 0.0001$, $p = 0.010$, $p = 0.0001$, $p = 0.0078$). Therefore, aging-induced reduction in the number of synapses, the length of synaptic arc strings, and the thickness of postsynaptic density, as well as the increase in protrusion gaps in hippocampal neurons could be reversed by aerobic exercise and intraperitoneal injection of exogenous irisin.

Exercise and irisin interventions optimized gut microbiota and metabolites in feces of aged mice

Exercise and irisin interventions rescued imbalanced gut microbiota of aged mice

Gut microbiota is an important mediator for reducing cognitive impairment during exercise [22], but whether exercise-induced irisin and optimized gut microbiota have the capability to suppress

cognitive impairment during aging process is still unknown. Therefore, in the present study, gut microbiota of aging-induced mice following exercise and irisin interventions were analyzed by 16S rRNA sequencing. Four animal experimental groups with 6 mice in each group, 1,527,506 original reads were obtained from a total of 24 samples, and 1,126,634 valid reads were obtained after quality control. After all samples were clustered according to 97% homology, 80,165 OTUs were obtained. The number of OTUs were 976, 640, 698 and 690 in YC, OC, OE, and OI groups, respectively (Fig. 4a), indicating aging-induced disturbance of gut microbiota, but exercise and irisin interventions can alleviate this disturbance. The species abundance at the top 10 phylum and genus levels showed that *Bacteroidetes*, *Firmicutes* and *Actinomycetes* were dominant at the phylum level (Fig. 4b), and *Ligilactobacillus* and *Lactobacillus Lachnospiraceae_NK4A136_group* were dominant at the genus level (Fig. 4c). The differential analysis of the top 10 species at the phylum and genus levels found that *Firmicutes/Bacteroidota* (F/B) ratio and *Actinobacteria* were significantly reduced in the OC group at the phylum level ($p = 0.0009$, $p = 0.0004$) when compared with the YC group (Fig. 4d and e); while significantly reversed this ratio and level in OI group ($p = 0.0302$, $p = 0.00024$), and the mice from the OE group with exercise intervention showed reversed *Actinobacteria* level ($p = 0.0094$). Similarly, compared with the YC group, the level of *Unidentified_Bacteria* in the OC group was significantly increased ($p = 0.0238$) (Fig. 4f), while the mice from the OE group exhibited a significantly reversed trend for this change ($p = 0.0262$), and the OI group showed a reversal sign, but no significant difference. At the genus level, *Bacteroides*, *Turicibacter*, *Parabacteroides* and *Bifidobacterium* in the OC group were reduced ($p = 0.0017$, $p = 0.0148$, $p = 0.0004$, $p = 0.0005$), while the OE and OI groups present a reverse in the reduction of *Bacteroides*, *Parabacteroides* and *Bifidobacterium* ($p = 0.0357$, $p = 0.0035$; $p = 0.0181$, $p = 0.0197$; $p = 0.0094$, $p = 0.0024$) (Fig. 4g-i and Supplementary Fig. S1), and the OE group did not exhibit the decline of *Turicibacter* (Fig. 4h) when compared with the YC group, suggesting that exercise and irisin interventions should participate in optimizing the balance of gut microbiota during aging process.

The principal coordinate analysis (PCoA) showed the distribution of bacterial community structure among different groups. ADONIS analysis showed the difference between YC and OC groups ($p = 0.001$), and between OC and OI groups ($p = 0.003$). However, the OC and OE groups have no difference ($p = 0.092$) (Fig. 4j). The Good Coverage rate of each sample was above 99%, and the α -diversity analysis (Fig. 4k-n) of gut microbiota from each group was carried out through observed species, chao1, ACE and Shannon index. The observed species, chao1, ACE and Shannon in the OC group were reduced significantly ($p = 0.0001$), and the mice from the OI group significantly reversed this trend ($p = 0.007$, $p = 0.0009$, $p = 0.0128$, $p = 0.0004$) when compared with the YC group. Although the mice from the OE group had a reversed trend, the statistical results were insignificant. These results suggested that intraperitoneal injection of irisin could reverse the change in the abundance and structure of gut microbiota caused by aging, thus contributing the compositions of gut microbiota like younger individuals, although exercise showed a particular improvement of recognitive capacity even without significant difference.


At the same time, potential biomarkers were screened through LEfSe analysis (LDA > 4) for differential microbiota in different groups (Fig. 4o). *Bacteroides*, *Bacterodaceae*, *Oscillospirales* and *Clostridia_UCG_014* were enriched in the YC group, *Ruminococcaceae* and *Oscillospirales* were enriched in the OC group, *Lleibacterium* and *Lleibacterium_valens* were enriched in the OE group, and *Ligilactobacillus*, *Lactobacillus_murinus*, *Actinobacteria*, *unidentified_Actinobacteria*, *Bifibobacteriales*, *Bifibobacterium*, *Bifibobacteriaceae* and *Prevotellaceae_UCG_001* were more abundant in the OI group.

In addition, KEGG functional prediction and cluster analysis of gut microbiota based on Tax4Fun found (Fig. 4q) that the aging of the mice could cause the changes in the compositions of gut microbiota, which may be an important factor for inducing and exacerbating the disease. Aerobic exercise and intraperitoneal injection of exogenous irisin may rescue the imbalanced gut microbiota to alleviate aging, inflammation, and neurodegeneration-related diseases.

Exercise and irisin interventions regulated fecal non-targeted metabolites in aged mice

Exercise in regulating fecal metabolites has been reported in many studies [23] and myokine irisin plays an active role in the regulation of gut microbiota. Therefore, quality control on fecal non-targeted metabolites was conducted. OPLS-DA analysis (Fig. 5a) showed that non-targeted metabolites could effectively distinguish the differential metabolites in feces of the mice from YC, OC, OE, and OI groups. In the statistical analysis of detected metabolites ($|\log_2 FC| \geq 1$, $VIP \geq 1$), compared with the YC group, 147 up-regulated and 173 down-regulated metabolites were detected in the OC group (Fig. 5b); compared with the OC group, 225 down-regulated and 64 up-regulated metabolites in the OE group were screened; similarly, 187 down-regulated and 45 up-regulated metabolites in the OI group were screened (Supplementary Fig. S2). Among them, compared with the YC group, 2-hydroxy-3,5-dinitro benzoic acid and 7-keto deoxycholic acid were decreased in mice from the OC group; while both exercise and irisin interventions could result in the significant increase of these compounds. Compared with the YC group, aging could induce the decrease of tetrahydro curcumin, while exercise intervention could reverse its decrease. Compared with the YC group, dihydrouracil and S-ribosyl-L-homocysteine in the OC group were reduced; while irisin intervention could rescue their reduction. Based on KEGG analysis, 94 pathways were enriched between YC and OC groups, 118 pathways were enriched between OC and OE groups, and 83 pathways were enriched between OC and OI groups. These pathways with significant differences were involved in serotonergic synapse ($p = 0.0197$) and cAMP signaling pathway ($p = 0.0197$) between YC and OC groups (Fig. 5c); pentose phosphate pathway ($p = 0.0052$), neuroactive ligand-receptor interaction ($p = 0.0138$), gap junction ($p = 0.0197$), synaptic vesicle cycle ($p = 0.0206$), taste transduction ($p = 0.0490$) between OC and OE groups (Fig. 5d). Gonadotropin-releasing hormone (GnRH) secretion ($p = 0.0341$) and cAMP signaling pathway ($p = 0.0463$) were predicted between OC and OI groups (Fig. 5e), which may be related to nervous system function.

To further explore the effect of exercise and irisin interventions on fecal microbiome and metabolites, a correlation analysis was conducted by extracting top 30 differential metabolites and microorganisms (Fig. 5f and Supplementary Table S1). The statistical difference was observed with the reduction of *Bifidobacterium*, *Turicibacter*, *Parabacteroides* and *Bacteroides* in mice from the OC group at the genus level, partially reversed in OE and OI groups (Fig. 4g-i and Supplementary Fig. S1). Among them, *Bifidobacterium* was significantly positively correlated with 2-(2R)-2-methyl-2-pyrrolidinyl-1H-benzimidazole-7-carboxamide, and pantothenate ($p = 0.0231$, $p = 0.0417$). *Parabacteroides* was significantly positively correlated with cyclamic acid, 2-(2R)-2-methyl-2-pyrrolidinyl-1H-benzimidazole-7-carboxamide, pantothenate, L-phenylalanine, and dodecanedioic acid ($p = 0.0010$, $p = 0.0345$, $p = 0.0172$, $p = 0.0351$, $p = 0.0001$). *Turicibacter* was significantly positively correlated with cyclamic acid ($p = 0.0009$). *Bacteroides* was significantly negatively correlated with sn-glycero-3-phosphocholine, methylcysteine and heparin ($p = 0.0093$, $p = 0.0212$, $p = 0.0246$), and positively correlated with cyclamic acid ($p = 0.0048$). These results suggest that exercise and irisin-induced changes in gut microbiota may also affect the compositions of metabolites in






Fig. 5. The analysis of non-targeted metabolites in feces of mice from each group and its correlation with gut microbiota. (a) OPLS-DA analysis; (b) The volcano plot of differential metabolites between YC and OC groups; (c-e) The functional KEGG analysis of differential metabolites; (f) The correlation analysis between differential gut microbiota and metabolites. Columns are microorganisms, red indicates a positive correlation and blue indicates a negative correlation, as well as the greater absolute value of the correlation labeled with darker color.

aging mice. Changes in metabolites may induce or exacerbate the progression of diseases. Therefore, modulating gut microbiota may be essential to alter metabolites to intervene diseases.

Exercise and irisin interventions regulated miRNAs and mRNAs in hippocampal tissues of aged mice

The changes of miRNAs in hippocampal tissues of the mice from YC and OE groups were evaluated by miRNA sequencing to uncover the regulatory roles of miRNAs. Quality control, statistical analysis, and the analysis of the original off-machine data ($|\log_2 FC| \geq 1$, $p \leq 0.05$) found that compared with the YC group, 19 down-regulated and 12 up-regulated miRNAs in hippocampal tissues of

the mice from the OC group were detected. Compared with the OC group, 7 down-regulated and 15 up-regulated miRNAs in the OE group, and 2 down-regulated and 8 up-regulated miRNAs in the OI group were detected, and a heatmap of the differential miRNAs in each group was established (Fig. 6a and Supplementary Fig. S3). The interaction network between partial miRNAs and genes in YC and OC groups was established to identify key miRNAs and corresponding target genes (Fig. 6b), and to list the significant differences in miRNAs and predicted target genes between each group (Supplementary Table S2-4). KEGG pathway enrichment of the predicted target genes of miRNAs in YC and OC groups was used to identify corresponding signal pathways, including ErbB,

Fig. 4. The 16S rRNA analysis and function prediction of gut microbiota in mice from each group. (a) The number of OTUs in mice from different groups; (b, c) The species compositions at the phylum and genus levels; (d-f) The TOP10 significantly different microorganisms at the phylum level in each group; (g-i) The TOP10 significantly different microorganisms at the genus level in each group; (j) PCA of fecal microorganisms in mice from each group; (k-n) Observed species, ACE, Shannon, and chao1; (o, p) The markers of LEfSe and differential gut microbiota; (d) Tax4Fun analysis and functional clustering of gut microbiota.

MAPK, cAMP, glutamatergic synapse, Toll-like receptor, Notch, Wnt, mTOR, Ras, apoptosis, insulin, mitophagy, NF- κ B, TNF- α , and other signal pathways, with the involvement of inflammation and aging-induced neurodegenerative diseases (Fig. 6c), and to list the enrichment of KEGG pathways for miRNAs and predicted target genes between each group (Supplementary Table S5-7).

Similarly, after extracting total RNA of hippocampal tissues, mRNA sequencing was also carried out to screen differential mRNAs ($|\log_2FC| \geq 1$, $p \leq 0.05$) between groups after statistical analysis of the effective data (transcripts per kilobase million, TPM). Compared with the YC group, 64 up-regulated and 28 down-regulated mRNAs in the OC group were detected. Compared with the OC group, 50 down-regulated and 11 up-regulated mRNAs in the OE group, and 79 down-regulated and 9 up-regulated mRNAs in the OI group were screened. The significantly differential genes were visualized (Fig. 6d, and Supplementary Table S8-10). Differential genes were enriched by KEGG (YC group vs. OC group), with major involvement in Toll-like receptor signal pathway, vascular smooth muscle contraction, dopaminergic synapse, NF- κ B, cAMP, B cell receptor, Ras, longevity regulation, T cell receptor, TNF- α , insulin signaling, MAPK and other signal pathways (Fig. 6e), and the KEGG pathway enrichment of each group (Supplementary Table S11-13). Joint analysis was performed based on miRNA and mRNA sequencing, and differentially enriched pathways between different groups were predicted after KEGG enrichment analysis (Fig. 6f). Among them, the opposite signal pathways enriched in YC and OC groups could be observed that the differential pathways were mainly involved in ErbB, Toll, neurotrophin, circadian rhythm, aminoacyl-tRNA biosynthesis, apoptosis, Fc receptor of immunoglobulin E (Fc epsilon RI), prolactin, GnRH, glycerophospholipid metabolism, estrogen, NF-kappa B, retrograde endocannabinoid signaling, T cell receptor, TNF, natural killer cell-mediated cytotoxicity, insulin, phospholipase D, Hippo, cell adhesion molecules (CAMs), protein processing in the endoplasmic reticulum, axon guidance, focal adhesion signaling pathway, actin cytoskeleton, Ras, and MAPK signal pathways. The enrichment of KEGG pathways for the joint analysis of miRNAs and mRNAs between each group were also listed (Supplementary Table S14-16).

From above results, Toll, inflammation, and insulin signal pathways may be the major mechanisms involved in the changes of hippocampal tissues from aged mice upon exercise and irisin interventions, suggesting the “gut-brain” crosstalk efficiency.

Correlation analysis between gut microbiota and miRNAs and mRNAs in hippocampal tissues of aged mice upon exercise and irisin interventions

To further explore the role of exercise and irisin interventions in the “gut-brain” axis, spearman correlation analysis for gut microbiota, miRNAs and mRNAs in hippocampal tissues between OC group and other groups were conducted to reveal a significant difference in *A2*, *Rikenella*, *Bifidobacterium*, *Parabacteroides*, *Monoglobus*, *Clostridium_sensu_stricto_1*, *Colidextribacter*, *Lachnolcostridium* and *Christensenellaceae_R-7* group at the genus level (Fig. 7a-b), and 46 differential miRNAs (Fig. 7c-d). Similarly, the correlation analysis between differential gut microbiota and 46 differential miRNAs was achieved (Fig. 7e). *A2*, *Rikenella*, *Bifidobacterium*, *Parabacteroides* and *Christensenellaceae_R-7* group were significantly positively correlated with miR-1264-5p, miR-1264-3p, miR-1298-5p, miR-1912-3p, miR-204-5p, miR-211-5p, miR-34c/b-5p, miR-448-3p, miR-467d-5p, miR-467c-5p, novel-695, miR-196a-5p, and negatively correlated with novel-72a, novel-43u, miR-1957a, novel-420, miR-206-3p, novel-428a/b, novel-43, novel-869a/b/c, miR-130b-3p, miR-141-3p, miR-1251-5p and

miR-200c-3p. *Lachnolcostridium* was positively correlated with miR-1264-5p, miR-1264-3p, miR-1298-5p, miR-1912-3p, miR-204-5p, miR-211-5p, miR-34c/b-5p, miR-448-3p, miR-1969, miR-1195, miR-196a/b-5p, novel-915, miR-467c/d-5p, novel-695, novel-95a/b/c/d, novel-67, miR-200b-5p, novel-101a/b, miR-10b-5p, miR-874a/b, novel-954, and significantly negatively correlated with novel-945, novel-730, novel-72a, novel-43u, miR-1957a, novel-420, miR-206-3p, novel-428a/b, novel-43, novel-869a/b/c, miR-130b-3p, miR-141-3p, miR-1251-5p, and miR-200c-3p. *Monoglobus* was positively correlated with miR-467d-5p, miR-467c-5p, novel-695 and miR-196a-5p, and negatively correlated with novel-72a, novel-43u and miR-1957a. *Clostridium_sensu_stricto_1* was negatively correlated with miR-34b-5p, novel-730 and miR-196a-5p, and positively correlated with novel-95b/c, novel-95a/c/d, miR-1957a, miR-1251-5p and miR-200c-3p. *Colidextribacter* was negatively correlated with miR-1264-5p, miR-1264-3p, miR-1298-5p, miR-1912-3p, miR-204-5p, miR-211-5p, miR-34c-5p, miR-448-3p, miR-34b-5p and novel-945, and positively correlated with novel-67, novel-101a, miR-141-3p, miR-1251-5p and miR-200c-3p. *Clostridium_sensu_stricto_1* was negatively correlated with miR-34b-3p, novel-730, miR-196a-5p, and positively correlated with novel-95a/b/c/d, miR-1957a, miR-141-3p, miR-1251-5p and miR-200c-3p.

At the same time, the differentially expressed gene transcripts in the OC group were screened and 747 genes were differentially expressed in the YC group, 681 genes in the OE group, and 1068 genes in the OI group; among them, there were 35 common genes (Fig. 7f-g), with correlation with gut microbiota in YC, OE, and OI groups, which revealed the significant difference from the OC group (Fig. 7h). It was found that *Parabacteroides* was positively correlated with *Cpne1*, *Pak3*, *Sema4a*, *Wdr48*, *Adra1a*, *Hspa8*, *Ly6e*, *Mtm1*, *Dpagt1*, *Hplbp3*, *Clvs1* and *Abcb8*, and negatively correlated with *Gbp6*, *U2af2*, *A930004D18Rik*, *Cxcr4*, *Ica1*, *Mecp2*, *Caln1*, *Hcfc2*, and *Scmh1*; *Bifidobacterium* was positively correlated with *Wdr48*, *Adra1a*, *Hspa8*, *Ly6e* and *Mtm1*, and negatively correlated with *Hcfc2* and *Prdm16*; *A2* was negatively correlated with *Clpb*, and positively correlated with *Tbc1d24*; *Rikenella* was negatively correlated with *Magi2* and *Akt1s1*, and positively correlated with *Mndal*; *Monoglobus*, *Christensenellaceae_R-7*, *Clostridium_sensu_stricto_1*, *Colidextribacter*, and *Lachnolcostridium* were positively correlated with *Cpne1*, *Pak3* and *Sema4a*, and negatively correlated with *Scmh1*. These results indicated that the correlation analysis of gut microbiota and miRNAs and mRNAs in hippocampal tissues of the mice from YC, OE, and OI groups was different from OC group, which may be the possible mechanisms for regulating gene expression in hippocampal tissues from the perspective of gut flora.

Exercise and irisin interventions alleviated inflammatory responses and TLR4/MyD88 signal pathway-mediated insulin resistance

From the analysis of fecal microbes, non-targeted metabolites, and screened miRNAs and mRNAs in hippocampal tissues, the screened signal pathways were predicted and enriched to be correlated with aging, inflammation, and neurodegenerative diseases. Toll receptor and insulin signal pathway were enriched with miRNAs and mRNAs through the analysis of KEGG pathways in hippocampal tissue. Lipopolysaccharide-binding protein (LBP) activating factor and IL-1 β , IL-6 and TNF- α from each group were detected. These results showed that LPS, LBP, IL-1 β , IL-6, and TNF- α in serum of the mice from the OC group were significantly increased ($p = 0.0405$, $p = 0.0287$, $p = 0.0089$, $p = 0.0188$, $p = 0.0149$) when compared with the YC group, while LPS, LBP, IL-1 β and TNF- α in serum of the mice from OE and OI groups were significantly decreased ($p = 0.0074$, $p = 0.0317$, $p = 0.0192$,




Fig. 6. Hippocampal miRNAs, mRNAs and joint analysis of the mice from each group. (a) Behavioral differential miRNAs, listed as groups; (b) Red indicates miRNAs and blue indicates target genes; (c) KEGG functional enrichment of differential miRNA target genes; (d) Heat map of differential genes in hippocampal tissues of the mice from each group. (e) KEGG functional enrichment of differential mRNAs; (f) The joint analysis of the difference between miRNAs and mRNAs. Red indicates a positive correlation, blue indicates a negative correlation, and darker color reveals higher correlation.




Fig. 7. The correlation analysis of gut microbiota and miRNAs and mRNAs in YC, OE, and OI groups differed from OC group. (a) The number of gut microbiota in YC, OE, and OI groups differed from OC group; (b) The differential microbiota in each group at the genus level; (c, d) The number and names of miRNAs in YC, OE, and OI groups differed from OC group; (e) The clustering heatmap of correlation between miRNAs and genus-level gut microbiota in YC, OE, and OI groups differed from OC group; (f) The number and names of differential genes in YC, OE, and OI groups differed from OC group; (g) The clustering heatmap of the correlation between genes and genus-level differential gut microbiota and genes. Red indicates a positive correlation, blue indicates a negative correlation, and darker indicates a more significant correlation.

$p = 0.0219$; $p = 0.0001$, $p = 0.0055$, $p = 0.0024$, $p = 0.0430$, $p = 0.0384$) (Fig. 8a). The upstream and downstream of TLR4-mediated inflammatory signal pathways were verified by qRT-PCR. *TLR4*, *MyD88*, *NF- κ B*, *IL-1 β* , *IL-6*, and *TNF- α* were significantly up-regulated in hippocampal tissues of the mice from the OC group ($p = 0.0039$, $p = 0.0084$, $p = 0.0254$, $p = 0.0010$, $p = 0.007$, $p = 0.0125$) when compared with the YC group; compared with the OC group,

TLR4, *MyD88*, *NF- κ B*, and *IL-1 β* were significantly decreased in the OE group ($p = 0.0019$, $p = 0.0424$, $p = 0.0153$, $p = 0.0056$, $p = 0.0006$), and *LBP*, *TLR4*, *MyD88*, *JNK*, and *IL-1 β* were significantly reduced in the OI group ($p = 0.0425$, $p = 0.0016$, $p = 0.0123$, $p = 0.0079$, $p = 0.0036$, $p = 0.001$) (Fig. 8b). Moreover, during detecting the inflammatory state of hippocampal tissue in each group were stained by Iba1, GFAP and p-NF- κ B p65, com-




Fig. 8. The serum inflammatory factors, hippocampal TLR4/MyD88 signal pathway-related genes, and corresponding proteins in hippocampal tissues of the mice from each group. (a) Serum levels of inflammation-related factors in different groups; (b) Hippocampal TLR4/MyD88 signal pathway-related inflammatory mRNA levels in each group; (c) Immunofluorescent staining of Iba1 and GFAP in hippocampal tissues of the mice from each group; (d) Corresponding inflammation levels in hippocampal tissues of the mice from each group evaluated by protein expression of p-NF- κ B p65 through immunofluorescence staining; (e) Statistical analysis for expression levels of corresponding inflammatory proteins in hippocampal tissues of the mice from each group.

pared with YC group, the activation levels of hippocampal microglial cells and NF- κ B p65 were increased in the OC group, but the activation of hippocampal microglial cells and NF- κ B p65 in OE and OI groups were decreased when compared with the OC group (Fig. 8c and e).

Furthermore, the expression of proteins associated with the inflammatory activation status was evaluated by Western blot. TLR4, MyD88, p-p38 MAPK/p38 MAPK, p-MKK7/MKK7, p-JNK/JNK, IL-1 β /6, TNF- α , p-IRS^{ser307}/IRS were significantly increased ($p = 0.0012$, $p = 0.0009$, $p = 0.0219$, $p = 0.0088$, $p = 0.0160$, $p = 0.001$, $p = 0.0046$, $p = 0.0305$, $p = 0.0005$) in the OC group when compared with the YC group (Fig. 8d and f). Compared with the OC group, TLR4, MyD88, IL-1 β and p-IRS^{ser307}/IRS were decreased ($p = 0.0287$, $p = 0.0027$, $p = 0.0264$, $p = 0.0263$) in the OE group, and TLR4, MyD88, p-p38 MAPK/p38 MAPK, p-MKK7/MKK7, p-JNK/JNK, IL-1 β /6 and p-IRS^{ser307}/IRS were significantly decreased ($p = 0.0144$, $p = 0.0253$, $p = 0.0126$, $p = 0.0119$, $p = 0.0298$, $p = 0.0006$, $p = 0.0069$, $p = 0.001$) in the OI group. The above results

indicated that hippocampal tissues of aged mice might induce neuronal inflammation and insulin resistance mediated by TLR4/MyD88 signal pathway through the activation of microglia by LPS. Exercise and irisin interventions could inhibit LPS in the body, alleviate the activation of microglial cells, and suppress hippocampal inflammation and insulin resistance.

Discussion

Exercise-modulating gut microbiota may be related to increasing irisin levels in the body. The sequencing of fecal 16S rRNA, non-targeted metabolites, and hippocampal miRNAs and mRNAs was conducted to further validate the effects of exercise and irisin-regulated gut microbiota and metabolites on miRNAs and genes in hippocampal tissues. The gut microbiota, genes, and signal pathways involved in these analyses may become potential intervention targets for improving cognitive capacity, among which, the activation of the TLR4 signal pathway in brain tissue mediated

by increased LPS due to the imbalanced gut microbiota may be an important regulator. Exercise suppresses neuroinflammation mediated by TLR4/MyD88 signal pathway in brain tissues to enhance cognitive capacity, which may be related to exercise-induced irisin in the body.

Aging can lead to impaired cognitive capacity, while drugs, exercise, and nutritional supplements are currently central interventions. To better understand the effect of exercise on aging-induced cognitive impairment, moderate-intensity aerobic and anaerobic training as possible within a week with a duration of 45–60 min for each training period reveals a significant improvement on cognitive performance in older adults over the age of 50 [24], and aerobic exercise is superior in cognitive improvement [25]. Therefore, 45-min aerobic treadmill running with 5 times a week and a training period of 3 consecutive months was provided for 21-month-old mice to further explore the molecular mechanisms of aerobic exercise-induced cognitive improvement during aging process in the present study. The improvement of learning and memory capacity upon exercise interventions may be correlated with increased function and number of neurons. Meanwhile, the observation with hippocampal HE, Nissl and NeuN staining, and TEM examination revealed the increased neuronal number, synaptic arc length, and PSD thickness of neuronal synapses after exercise intervention, and the synaptic gap was decreased significantly [26], including the critical role of gut microbiota [27].

There is a significant difference in irisin level across different models. Regression analysis of disease states shows that plasma irisin level is positively correlated with normal and mildly impaired cognitive capacity of patients without AD. In contrast, higher irisin level in AD patients may have smaller hippocampal volume [28], indicating that the occurrence of the disease may be a feedback response from the increased irisin to slow down the disease progression and the depletion of irisin secretion when the disease state is irreversible. Relevant studies have also confirmed that exercise can enhance cognitive capacity due to the optimized gut microbiota [29,30]. In the present study, serum irisin levels in mice subjected to exercise and irisin interventions are significantly higher than the aging model mice, which is consistent with previous results [31]. Moreover, the direct injection of recombinant irisin has the stronger effect than exercise intervention. However, the studies on irisin-mediated optimization of gut microbiota are limited, and the regulation of gut microbiota by irisin is only reported in colitis and ischemia–reperfusion injury models [11,13]. It is worth noting that although exercise does not show a significantly regulatory effect on gut microbiota, it still shows a significantly reversal cognitive capacity during aging process, suggesting exercise-induced multiple benefits. Similarly, exercise can effectively regulate gut microbiota to prevent gut barrier damage [32]. However, the studies in mouse aging models are still limited. The limited regulation capacity to gut microbiota may be associated with lower exercise capacity and excessive oxidative stress in older mice [33,34]. In contrast, excessive exercise intensity may impact the structure and function of gut microbiota in the elderly population [35]. Meanwhile, some studies have also shown that exercise can reshape gut microbiota although the reshaping capacity is limited [36]. In the present study, exercise has a regulatory effect on gut microbiota, and is weaker than irisin intervention, which may be due to irisin circulation during exercise intervention when compared with direct irisin intervention. These results suggest that moderate-intensity aerobic exercise has a limited role in regulating gut microbiota in aged mice or exercise-induced irisin does not reach up to the level for triggering gut microbiota.

Our 16S RNA sequencing data of gut microbiota have confirmed that aging can lead to a significant decrease in *F/B* ratio and abundance of *Actinobacteria*, *unidentified_Bacteria*, *Parabacteroides*, *Bac-*

teroides, *Turicibacter*, and *Bifidobacterium*, whereas exercise and irisin injection can reverse the decline in the abundance of gut microbiota that is closely related to cognitive capacity. Statistical analysis of gut microbiota at the phylum and genus levels has documented that *F/B* ratio shows a significant decrease in aged mice, with consistent results with previous studies [37,38], and the *F/B* ratio decline may be also due to aging stages, and races [39–41]. In a previous study on gut microbiota and cognition in neurologically healthy elderly, it has found that *Firmicutes* is positively correlated with memory capacity, but *Bacteroidetes* is negatively correlated [42]. The fecal microbiota test of Amyotrophic lateral sclerosis patients with cognitive impairment and Amyotrophic lateral sclerosis patients with normal cognition also shows a significant increase in *F/B* ratio [43], which is consistent with the decrease in *F/B* ratio observed during aging process. In rats with memory impairment, the traditional Chinese herbal prescription Dangshen Yuanzhi Powder also can reverse the decrease in *F/B* ratio while improving memory capacity [44]. In the present study, exercise and irisin interventions reversed this ratio and showed the improved cognitive capacity, further confirming that *F/B* ratio may be a potential factor of cognitive capacity. However, in obesity models with cognitive impairment caused by diabetes, *F/B* ratio shows a significant increase [45,46], which may be related to unnatural aging and exogenous food or drug intervention. A significant increase in *Actinobacteria* has been found in both AD and elderly cognitive impairment populations [47–49]. However, *Actinobacteria* reveals an significant increase in studies using Triphala to improve cognition in AD mice [50], and *Actinobacteria* was a significant decrease in abundance with increasing age [51], which is also consistent with our results. In addition, the studies on the relationship between intestinal microbial changes and cognition in AD people have also confirmed that the abundance of *Actinobacteria* is significantly reduced [52]. Significant reduction in *Actinobacteria* has been found in 5XFAD mice with severe cognitive decline when compared with wild-type mice at the same age [53]. In another study, *Actinobacteria* is directly linked to cognitive capacity [54]. The traditional Chinese herb Jiedu Yizhi prescription can significantly improve cognitive capacity of APP/PS1 mice, and can also suppress the TLR4/NF- κ B signal pathway in the brain through inhibiting *Actinobacteria*, further suggesting that *Actinobacteria* may be involved in regulating signal pathways for controlling cognitive capacity [55]. In chronic neuropathic pain models, a selective decrease in gut flora accompanied by an increase in *Actinobacteria* may contribute to the occurrence of cognitive impairment [56]. PD also shows a rapid decline in cognitive level during the development process of the disease. A 3-year follow-up study has found that the reduction of *Actinobacteria* is closely related to accelerated cognitive deterioration [57]. *Actinobacteria* abundance also shows a significant increase when administrated with the cognitively beneficial Enoki mushroom polysaccharide [58]. However, *Actinobacteria* in the studies of aging and cognition fields has received attention only in recent years [59]. Although there is some controversy, more studies on its beneficial cognitive improvement may be needed to further confirm.

Based on our study, *Parabacteroides* reveals a significant decrease during aging process, while exercise and irisin interventions can reverse its aging-induced reduction. Similarly, a significant decrease in the abundance of *Parabacteroides* has also found in the elderly with mild cognitive impairment [60]. In hemodialysis patients with cognitive impairment, there is also a positive correlation between *Parabacteroides* and cognitive capacity [61]. Cognitive capacity can be enhanced by optimizing the compositions of gut flora upon probiotics supplementation, in which the abundance of *Parabacteroides* also shows a significant increase [62]. However, *Parabacteroides* show an increase in models of cognitive impairment after schizophrenia, stroke, and radiation-

induced cancers [63–65]. These opposite results may be related to different disease models, and the specific factors may require more in-depth study. The reduction of *Bacteroides* during aging process may be negatively correlated with body inflammation [66], and supplementing *Bacteroides* can inhibit inflammation levels [67]. Therefore, *Bacteroides* should have close association with cognitive improvement in the elderly [68], is positively correlated with cognitive capacity [69], as well as involved in neurodevelopment to improve infant cognitive capacity [70]. In clinical studies, fecal transplants can suppress cognitive decline due to the significant increase in *Bacteroides* [71]. Mendelian analysis of over 200,000 cognitive data and nearly 20,000 gut microbes has demonstrated that *Bacteroides* plays an active role in cognitive protection [72]. Some studies have shown a significant increase in *Bacteroides* by transplanting gut flora of AD persons into germ-free mice [73]. The detection of microbiota in people with dementia has also confirmed an increase in the abundance of *Bacteroides* as an independent factor in dementia [74].

In addition, the beneficial *Turicibacter* is also significantly reduced in aging mice, while exercise and irisin interventions rescue this deleterious reduction. Supplementing *Turicibacter* could delay aging process to a certain extent [75], and a significant decrease in *Turicibacter* abundance is also observed in APP/PS1 mice with poor cognitive capacity [76]. Dietary methionine restriction as a diet intervention for anti-aging and suppressing aging-induced cognitive impairment can stimulate cognitive capacity through significantly increasing the abundance of *Turicibacter* [77]. *Bifidobacterium* exhibits anti-aging effects on promoting cognitive capacity and metabolic level of AD mice [78,79], which is possibly correlated with neuron regeneration [28]. In a randomized controlled trial on the elderly, it is found that exogenous supplementation of *Bifidobacterium* can significantly improve the learning and memory capacity of elderly population [80]. Pregnant mice supplemented with omega-3 exhibit a significant increase of *Bifidobacterium* and enhanced cognitive capacity in their offsprings [81]. Aged mice and humans supplemented with *Bifidobacterium* have also found that cognitive function is significantly enhanced, which may be related to the improvement in the gut environment and barrier and inflammation levels [82,83]. Moreover, middle-aged mice supplemented with *Bifidobacterium* show improved cognitive capacity and metabolic function, further confirming its impact on brain function [84,85]. Supplementing *Bifidobacterium* has shown to modulate gut flora when rescuing mild cognitive impairment, with the involvement in improving synaptic plasticity through up-regulating brain-derived neurotrophic factor (BDNF), fibronectin type III domain-containing protein 5 (FNDC5), and postsynaptic density protein 95 (PSD-95) in the brain [86]. After exercise and irisin interventions in the present study, the level of *Bifidobacterium* in aged mice reveals a significant increase accompanied by the improvement of brain structure and cognitive capacity, further suggesting that exercise or exercise-induced irisin may promote the optimal balance in the abundance of beneficial bacteria.

The change in gut microbiota can induce corresponding change of the metabolites in feces. The fecal metabolites have significant differences in the OPLS-DA scores of the metabolites from different interventions. *Bacteroides*, *Turicibacter*, and *Bifidobacterium* at the genus level are significantly reduced in feces of aged mice, which can be reversed by exercise and irisin interventions. *Bifidobacterium* is significantly positively correlated with 1H-2-(2R)-2-methyl-2-pyrrolidinyl-benzimidazole-7-carboxamide, and pantothenate, among which 1H-2-(2R)-2-methyl-2-pyrrolidinyl-benzimidazole-7-carboxamide has limited research, while pantothenic acid is involved in improved cognitive capacity. Pantothenic acid is also known as vitamin B5, and the deficient intake of pantothenic acid may lead to cognitive decline [87]. In the analysis of human

cerebrospinal fluid components, it is found that pantothenic acid is significantly positively correlated with aging [88], and abnormal change in pantothenic acid is also found in gut flora of normal people and people transitioning from MCI to dementia [89]. *Turicibacter* is significantly positively correlated with cyclamic acid. However, the studies on the role of cyclamic acid in cognition seem to be lacking. *Bacteroides* is significantly negatively correlated with sn-glycero-3-phosphocholine, methylcysteine and heparin, and positively correlated with cyclamic acid. Sn-glycero-3-phosphocholine (synonym choline alfoscerate) as an acetylcholine precursor in improving cognition has been confirmed by many studies. Sn-glycero-3-phosphocholine has a positive effect on cognitive improvement in natural aging and different stages of AD populations [90,91], and is even considered a cognitive enhancer. A meta-analysis involving 449 population-based trials has demonstrated that phosphocholine supplementation can significantly improve cognition [92]. However, the relationship between methylcysteine, heparin, and cyclohexanoic acid and cognition has been less studied.

To explore the relationship between gut microbiota and miRNAs and mRNAs in hippocampal tissues, differential gut microbiota between YC, OE, OI, and OC groups at the genus level are screened. Among differential miRNAs, miR-206-3p is significantly up-regulated in the OC group and down-regulated in YC, OC and OI groups at a certain degree. The miR-206-3p is directly involved in regulating BDNF [93], which may become a potential marker for diagnostic and treatment tools for AD [94]. In addition, the relative expression of miR-103-3p and miR-103b-3p was higher in OC and OE groups. Relevant studies have shown that lower levels of miR-103-3p in human serum are independently associated with cognitive impairment [95]. Based on sequencing data of AD patients and normal populations, miR-103-3p can interact with most of the key genes screened to predict AD [96]. MiR-103-3p in the OE group seems to be consistent with that in the OC group, but relevant studies have confirmed that exercise may cause the increase of miR-103-3p to a certain extent [97], but the studies between miR-103b-3p and cognitive capacity have not been reported. In transcriptome sequencing, we have found that some genes closely correlated with cognition are significantly different between YC and OC groups, such as *Cst7*, *Klk6*, *Slc13a4*, *Trem2*, *Slc16a9*, *Cldn1*, and *Apod*. The *Cst7* gene is up-regulated in the OC group, but down-regulated in the YC and OI groups, and is also associated with cognitive decline [98], while the inhibition of *Cst7* shows cognitive deficits [99]. As a gene closely related to AD, the expression of *Klk6* can lead to neuronal degeneration [100], and *Klk6* has been found to be significantly elevated in plasma of AD persons, and may become a biomarker for predicting the severity of dementia AD [101]. *Slc13a4* has sodium-sulfate symporter activity, and the deletion of the *Slc13a4* gene causes a significant decrease in the learning and memory capacity of mice, but in contrast, *Slc13a4*^{+/-} mice show an increase in the appearance of hippocampal neurons [102]. *Trem2* has been found to be related to AD in many studies and is highly expressed in AD. Inhibiting *Trem2* has also become an important target to alleviate cognitive decline [103,104]. *Slc16a9* methylation mediates the effects of prenatal alcohol exposure on neonatal cognitive and attention-related deficits [105]. In the blood-brain barrier endothelial cells of aged mice, TLR4 activation is accompanied by a decrease in *Cldn1*, while the downregulation of TLR4 shows an increase in *Cldn1*, and *Cldn1* is positively correlated with cognition [106]. Leukoaraiosis patients with cognitive impairment also show a significant decrease in *Cldn1* [107]. Moreover, the inhibition of *ACE* and *ACE* heterozygous mice show an exacerbation of AD symptoms [108], and the underlying mechanism may be related to the degradation of β -amyloid protein [109]. However, there are many studies showing that *ACE* is not correlated with cognition [110], or that blocking *ACE* can

alleviate further deterioration of cognition [111]. Therefore, the relationship between *ACE* and cognition may need further exploration. *Apod* shows the up-regulation with the extension of age [112], also shows high expression in AD patients [113] and a negative correlation with cognitive capacity [114]. *Gm28294* and *C5ar2* are significantly down-regulated in the OC group and increased in YC, OE and OI groups. Unfortunately, there are no reports on *Gm28294* in the literature. *C5ar2* may play a neuroprotective role in AD and delay the occurrence of AD, complement and astrocyte-related genes to a certain extent [115]. *Gpr17* and *Fos* are significantly down-regulated in the OC group and up-regulated in the OE group. However, some studies have shown that knocking out and inhibiting *Gpr17* exhibits positive effects on cognition [116,117], which is contrary to our results. *Fos* gene plays an important role in spatial memory [118]. The c-Fos as one of the *Fos* gene members is dysregulated in AD patients [119], and is also significantly down-regulated in aging mice with cognitive deficits [120], while running training can significantly up-regulate its expression in hippocampal tissues [121], which is consistent with our results.

U2af2, *Gbp6*, *Scmh1*, *Prdm16*, *Akt1s1*, *Caln1*, *Mecp2*, *Cxcr4*, *A930004D18Rik*, *Ica1*, *Hcfc2*, *Prdm16* and *Magi2* genes are up-regulated in the OC group, while *Adra1a*, *Glmp*, *Ctge5*, *Ndel1*, *Wdr48*, *Tenm4*, *Hspa8*, *Pak3*, *Clvs1*, *Cpne1*, *Sema4a*, *Ly6e*, *Abcb8*, *Dpagt1* and *Hp1bp3* are down-regulated in the OC group, which may be the important predictors of aging. Based on the correlation analysis of gut microbiota *A2*, *Rikenella*, *Bifidobacterium*, *Parabacteroides*, *Monoglobus*, *Christensenellaceae_R-7*, *Clostridium_sensu_stricto_1*, *Colidextribacter* and *Lachnolcostridium* at the genus level and miRNAs and mRNAs, gut microbiota has a significant potential role in influencing hippocampal miRNAs and mRNAs to participate in aging-induced neurological diseases (Fig. 6h).

Inflammation-induced insulin resistance is one of the major inducers of aging-induced cognitive impairment, which may be closely correlated with the increased LPS and LBP in the body. Both exercise and irisin interventions have been confirmed to exert anti-inflammatory effects with the focus on TLR4/MyD88 [14,122,123]. Under normal physiological conditions, LPS in gut microbiota can maintain homeostasis at a certain level. At aging and disease states, abnormal changes in gut microbiota may lead to the release of a large amount of LPS, thereby aggravating the inflammatory response of the whole body and organs, and even inducing insulin resistance and the occurrence of diseases. TLR4 signal pathway is the primary inflammatory signal for LPS-induced LBP activation. In the present study, suppressing TLR4 signal pathway in hippocampal tissues of aged mice upon exercise and irisin interventions can suppress inflammation. Similarly, the significant up-regulation of *TLR4*, *MYD88*, *IL-1 β* , and *IL-6* genes in hippocampal tissues of the elderly has also been confirmed by human experiments [124,125]. TLR4 may mediate the transduction of IL-1 β signal pathway in hippocampal tissues of aged mice. The absence of TLR4 can offer aged mice for better learning and memory functions [20,126], which may be closely correlated with the suppression of inflammation and the mitigation of neuronal disorders [127], illustrating that the activation of TLR4 may be one of the important causes of insulin resistance and accelerating cognitive impairment during aging process [128]. During natural aging or drug-induced aging process, p-p38 MAPK in hippocampal tissues can be triggered to a certain extent [129,130], and aerobic exercise can significantly result in its down-regulation [131], which is consistent with our study, suggesting an obviously positive anti-inflammation efficiency and enhanced insulin sensitivity upon exercise and irisin interventions. However, the effect of irisin intervention on inhibiting hippocampal neuroinflammation and insulin resistance is better than that of exercise intervention, which may be related to the fact that the irisin level in aging mice induced

by moderate exercise is not enough to optimize gut microbiota for suppressing inflammation. The specific mechanism needs to be further clarified.

In the present study, comprehensive testing of gut flora, non-targeted metabolites, hippocampal miRNAs and mRNAs, blood-related indicators, and hippocampal tissue samples shows that irisin intervention is more effective than exercise. Exercise-induced regulation of the gut-brain axis may be closely related to resultant endocrine, metabolite and physiological changes involved in regulating gut flora. Therefore, exercise-induced irisin can modulate gut microbiota to improve cognition, which is also a new way to uncover exercise to improve cognition. Analyzing how exercise regulates gut flora and participates in disease improvement may be a focus of future research. At present, it has been confirmed that irisin is positively correlated with cognitive capacity of AD persons with cognitive impairment [132], and exercise plays a role in enhancing cognition by increasing irisin levels [13]. However, exercise seems to be a reference only for people with normal physical activities, but not for people with cognitive impairment, physical disabilities or those who are unable to accomplish exercise. Therefore, direct intervention with irisin to achieve the mimic effects of exercise on health promotion and disease prevention or rehabilitation is the ultimate goal that we are currently pursuing. Unfortunately, there are currently no experimental reports on irisin interventions in humans, but exercise [13], natto kinase [133], low-intensity pulse [134], solanum melongena extract [135] can trigger the circulating level of irisin in the body. Therefore, increasing the circulating level of irisin in the body through exogenous interventions targeting the body's irisin may also be a potential strategy for disease interventions in the future. It is worth noting that irisin appears to vary in different diseases, and if used in human treatment, more detailed screening of body functions and diseases may be required. The collection of data from larger samples of people with different ages and diseases and the exploration of human-related experiments may be a breakthrough in promoting the clinical value of irisin.

Conclusion

Exercise-reversed cognitive impairment of aging mice may be correlated with the generation and secretion of exercise-induced irisin and the irisin-mediated optimization of gut microbiota and metabolites through suppressing TLR4/MyD88 signal pathway and insulin resistance. This finding may provide a new reference for exercise-mediated gut microbiota to enhance cognitive capacity during aging process. Unfortunately, the beneficial effect and underlying mechanisms of irisin on optimizing the balance of gut microbiota are not fully validated. Moreover, irisin gene knockout to affect the regulation of gut microbiota and cognitive capacity may need to be further verified.

Compliance with Ethics Requirements

All experiments involving animals were conducted according to the ethical policies and procedures approved by Institutional Animal Care and Use Committee of Wuhan Sports University with the number of S087-21-05D, and all experiments involving animals were complied with the internationally recognized 3R principles.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (No. 32071176), the 14th Five-Year-Plan Advantageous and Characteristic Disciplines (Groups) of Colleges and Universities in Hubei Province for Exercise and Brain Science from Hubei Provincial Department of Education and the Leading Talent Program and Innovative Start-Up Foundation from Wuhan Sports University to NC.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jare.2024.10.031>.

References

- [1] Rosenberg PB, Oh ES. Association of subjective cognitive decline with postoperative complications could herald dementia risk. *Am J Geriatr Psychiatry* 2021;29(4):362–1334. doi: <https://doi.org/10.1016/j.jagp.2020.09.013>.
- [2] Desai R, Whitfield T, Said G, John A, Saunders R, Marchant NL, et al. Affective symptoms and risk of progression to mild cognitive impairment or dementia in subjective cognitive decline: A systematic review and meta-analysis. *Ageing Res Rev* 2021;71:101419. doi: <https://doi.org/10.1016/j.arr.2021.101419>.
- [3] Walker KA, Gottesman RF, Wu A, Knopman DS, Gross AL, Mosley TH, et al. Systemic inflammation during midlife and cognitive change over 20 years: The ARIC Study. *Neurology* 2019;92(11):e1256–67. doi: <https://doi.org/10.1212/wnl.00000000000007094>.
- [4] Glatigny M, Moriceau S, Rivagorda M, Ramos-Brossier M, Nascimbeni AC, Lante F, et al. Autophagy is required for memory formation and reverses age-related memory decline. *Curr Biol* 2019;29(3). doi: <https://doi.org/10.1016/j.cub.2018.12.021>. 435–48.e8.
- [5] Cabral-Miranda F, Tamburini G, Martinez G, Ardiles AO, Medinas DB, Gerakis Y, et al. Unfolded protein response IRE1/XBP1 signaling is required for healthy mammalian brain aging. *Embo j* 2022:e111952. doi: <https://doi.org/10.15252/emboj.202111952>.
- [6] Yaffe K, Falvey CM, Hoang T. Connections between sleep and cognition in older adults. *Lancet Neurol* 2014;13(10):1017–28. doi: [https://doi.org/10.1016/s1474-4422\(14\)70172-3](https://doi.org/10.1016/s1474-4422(14)70172-3).
- [7] O'Toole PW, Jeffery IB. Gut microbiota and aging. *Science* 2015;350(6265):1214–5. doi: <https://doi.org/10.1126/science.aac8469>.
- [8] Bialecka-Debek A, Granda D, Szmidt MK, Zielińska D. Gut Microbiota, probiotic interventions, and cognitive function in the elderly: A review of current knowledge. *Nutrients* 2021;13(8). doi: <https://doi.org/10.3390/nu13082514>.
- [9] Cheng LH, Chou PY, Hou AT, Huang CL, Shiu WL, Wang S. Lactobacillus paracasei PS23 improves cognitive deficits via modulating the hippocampal gene expression and the gut microbiota in D-galactose-induced aging mice. *Food Funct* 2022;13(9):5240–51. doi: <https://doi.org/10.1039/d2fo00165a>.
- [10] Ghosh TS, Rampelli S, Jeffery IB, Santoro A, Neto M, Capri M, et al. Mediterranean diet intervention alters the gut microbiome in older people reducing frailty and improving health status: the NU-AGE 1-year dietary intervention across five European countries. *Gut* 2020;69(7):1218–28. doi: <https://doi.org/10.1136/gutjnl-2019-319654>.
- [11] Huangfu LX, Cai XT, Yang JN, Wang HC, Li YX, Dai ZF, et al. Irisin attenuates inflammation in a mouse model of ulcerative colitis by altering the intestinal microbiota. *Exp Ther Med* 2021;22(6):1433. doi: <https://doi.org/10.3892/etm.2021.10868>.
- [12] Liu Q, Zhu Y, Li G, Guo T, Jin M, Xi D, et al. Irisin ameliorates myocardial ischemia-reperfusion injury by modulating gut microbiota and intestinal permeability in rats. *PLoS One* 2023;18(9):e0291022.
- [13] Islam MR, Valaris S, Young MF, Haley EB, Luo R, Bond SF, et al. Exercise hormone irisin is a critical regulator of cognitive function. *Nat Metab* 2021;3(8):1058–70. doi: <https://doi.org/10.1038/s42255-021-00438-z>.
- [14] Mazur-Bialy AI, Pocheć E, Zarawski M. Anti-inflammatory properties of irisin, mediator of physical activity, are connected with TLR4/MyD88 signaling pathway activation. *Int J Mol Sci* 2017;18(4):701. doi: <https://doi.org/10.3390/ijms18040701>.
- [15] Chidambaram SB, Essa MM, Rathipriya AG, Bishir M, Ray B, Mahalakshmi AM, et al. Gut dysbiosis, defective autophagy and altered immune responses in neurodegenerative diseases: Tales of a vicious cycle. *Pharmacol Ther* 2022;231:107988. DOI: 10.1016/j.pharmthera.2021.107988.
- [16] Scherer T, Sakamoto K, Buettner C. Brain insulin signalling in metabolic homeostasis and disease. *Nat Rev Endocrinol* 2021;17(8):468–83. doi: <https://doi.org/10.1038/s41574-021-00498-x>.
- [17] Rubio C, Lizárraga E, Álvarez-Cilleros D, Pérez-Pardo P, Sanmartín-Salinas P, Toledo-Lobo MV, et al. Aging in male wistar rats associates with changes in intestinal microbiota, gut structure, and cholecystokinin-mediated gut-brain axis function. *J Gerontol A Biol Sci Med Sci* 2021;76(11):1915–21. doi: <https://doi.org/10.1093/gerona/glaa313>.
- [18] Sun Y, Wang Y, Lin Z, Zhang F, Zhang Y, Ren T, et al. Irisin delays the onset of type 1 diabetes in NOD mice by enhancing intestinal barrier. *Int J Biol Macromol* 2024;130857. doi: <https://doi.org/10.1016/j.ijbiomac.2024.130857>.
- [19] Brandt A, Kromm F, Hernández-Arriaga A, Martínez Sánchez I, Bozkir H, Staltner R, et al. Cognitive alterations in old mice are associated with intestinal barrier dysfunction and induced toll-like receptor 2 and 4 signaling in different brain regions. *Cells* 2023;12(17):2153. doi: <https://doi.org/10.3390/cells12172153>.
- [20] Fei X, Dou YN, Lv W, Ding B, Wei J, Wu X, et al. TLR4 deletion improves cognitive brain function and structure in aged mice. *Neuroscience* 2022;4921. doi: <https://doi.org/10.1016/j.neuroscience.2022.04.007>.
- [21] Chen D, Zhang Y, Zhang M, Chang J, Zeng Z, Kou X, et al. Exercise attenuates brain aging by rescuing down-regulated wnt/ β -catenin signaling in aged rats. *Front Aging Neurosci* 2020;12:105. doi: <https://doi.org/10.3389/fnagi.2020.00105>.
- [22] Koblinsky ND, Power KA, Middleton L, Ferland G, Anderson ND. The role of the gut microbiome in diet and exercise effects on cognition: A review of the intervention literature. *J Gerontol A Biol Sci Med Sci* 2022;78(2):195. doi: <https://doi.org/10.1093/gerona/glab166>.
- [23] Qiu L, Gong F, Wu J, You D, Zhao Y, Xu L, et al. Exercise interventions improved sleep quality through regulating intestinal microbiota composition. *Int J Environ Res Public Health* 2022;19(19):12385. doi: <https://doi.org/10.3390/ijerph191912385>.
- [24] Northey JM, Cherbuin N, Pumpa KL, Smee DJ, Rattray B. Exercise interventions for cognitive function in adults older than 50: A systematic review with meta-analysis. *Br J Sports Med* 2018;52(3):154–60. doi: <https://doi.org/10.1136/bjsports-2016-096587>.
- [25] López-Ortiz S, Valenzuela PL, Seisdedos MM, Morales JS, Vega T, Castillo-García A, et al. Exercise interventions in Alzheimer's disease: A systematic review and meta-analysis of randomized controlled trials. *Ageing Res Rev* 2021;72:101479. doi: <https://doi.org/10.1016/j.arr.2021.101479>.
- [26] Siette J, Westbrook RF, Cotman C, Sidhu K, Zhu W, Sachdev P, et al. Age-specific effects of voluntary exercise on memory and the older brain. *Biol Psychiatry* 2013;73(5):435–42. doi: <https://doi.org/10.1016/j.biopsych.2012.05.034>.
- [27] Xu TC, Lv Y, Liu QY, Chen HS. Long-term atorvastatin improves cognitive decline by regulating gut function in naturally ageing rats. *Immun Ageing* 2022;19(1):52. doi: <https://doi.org/10.1186/s12979-022-00311-x>.
- [28] Kim H, Shin J, Kim S, Kim S, Cho B, Park SJ, et al. Bifidobacterium bifidum BGN4 and Bifidobacterium longum BORI promotes neuronal rejuvenation in aged mice. *Biochem Biophys Res Commun* 2022;60341–6038. doi: <https://doi.org/10.1016/j.bbrc.2022.03.024>.
- [29] Lai Z, Shan W, Li J, Min J, Zeng X, Zuo Z. Appropriate exercise level attenuates gut dysbiosis and valeric acid increase to improve neuroplasticity and cognitive function after surgery in mice. *Mol Psychiatry* 2021;26(12):7167–87. doi: <https://doi.org/10.1038/s41380-021-01291-y>.
- [30] Cryan JF, O'Riordan KJ, Sandhu K, Peterson V, Dinan TG. The gut microbiome in neurological disorders. *Lancet Neurol* 2020;19(2):179–94. doi: [https://doi.org/10.1016/s1474-4422\(19\)30356-4](https://doi.org/10.1016/s1474-4422(19)30356-4).
- [31] Hu C, Zhang X, Hu M, Teng T, Yuan YP, Song P, et al. Fibronectin type III domain-containing 5 improves aging-related cardiac dysfunction in mice. *Aging Cell* 2022;21(3):e13556.
- [32] Xia WJ, Xu ML, Yu XJ, Du MM, Li XH, Yang T, et al. Antihypertensive effects of exercise involve reshaping of gut microbiota and improvement of gut-brain axis in spontaneously hypertensive rat. *Gut Microbes* 2021;13(1):1–24. doi: <https://doi.org/10.1080/19490976.2020.1854642>.
- [33] Clark A, Mach N. Exercise-induced stress behavior, gut-microbiota-brain axis and diet: a systematic review for athletes. *J Int Soc Sports Nutr* 2016;1343. doi: <https://doi.org/10.1186/s12970-016-0155-6>.
- [34] Cook MD, Martin SA, Williams C, Whitlock K, Wallig MA, Pence BD, et al. Forced treadmill exercise training exacerbates inflammation and causes mortality while voluntary wheel training is protective in a mouse model of colitis. *Brain Behav Immun* 2013;3346–56. doi: <https://doi.org/10.1016/j.bbi.2013.05.005>.
- [35] Zhu Q, Jiang S, Du G. Effects of exercise frequency on the gut microbiota in elderly individuals. *Microbiologopen* 2020;9(8):e1053.
- [36] Lamoureux EV, Grandy SA, Langille MGI. Moderate exercise has limited but distinguishable effects on the mouse microbiome. *mSystems* 2017;2(4):e00066–e17. doi: <https://doi.org/10.1128/mSystems.00006-17>.
- [37] Vemuri R, Sherrill C, Davis MA, Kavanagh K. Age-related colonic mucosal microbiome community shifts in monkeys. *J Gerontol A Biol Sci Med Sci* 2021;76(11):1906–14. doi: <https://doi.org/10.1093/gerona/glaa256>.
- [38] Haran JP, McCormick BA. Aging, frailty, and the microbiome-how dysbiosis influences human aging and disease. *Gastroenterology* 2021;160(2):507–23. doi: <https://doi.org/10.1053/j.gastro.2020.09.060>.
- [39] Mariat D, Firmesse O, Levenez F, Guimaraes V, Sokol H, Doré J, et al. The Firmicutes/Bacteroidetes ratio of the human microbiota changes with age. *BMC Microbiol* 2009;9:123. doi: <https://doi.org/10.1186/1471-2180-9-123>.
- [40] Sychala MS, Venna VR, Jandzinski M, Doran SJ, Durgan DJ, Ganesh BP, et al. Age-related changes in the gut microbiota influence systemic inflammation and stroke outcome. *Ann Neurol* 2018;84(1):23–36. doi: <https://doi.org/10.1002/ana.25250>.

- [41] Vaiserman A, Romanenko M, Piven L, Moseiko V, Lushchak O, Kryzhanovska N, et al. Differences in the gut Firmicutes to Bacteroidetes ratio across age groups in healthy Ukrainian population. *BMC Microbiol* 2020;20(1):221. doi: <https://doi.org/10.1186/s12866-020-01903-7>.
- [42] Manderino L, Carroll I, Azcarate-Peril MA, Rochette A, Heinberg L, Peat C, et al. Preliminary evidence for an association between the composition of the gut microbiome and cognitive function in neurologically healthy older adults. *J Int Neuropsychol Soc* 2017;23(8):700–75. doi: <https://doi.org/10.1017/s155617717000492>.
- [43] Gong Z, Ba L, Tang J, Yang Y, Li Z, Liu M, et al. Gut microbiota links with cognitive impairment in amyotrophic lateral sclerosis: A multi-omics study. *J Biomed Res* 2022;37(2):125–37. doi: <https://doi.org/10.7555/jbr.36.20220198>.
- [44] Ren H, Gao S, Wang S, Wang J, Cheng Y, Wang Y, et al. Effects of Dangshen Yuanzhi Powder on learning ability and gut microflora in rats with memory disorder. *J Ethnopharmacol* 2022;296:115410. doi: <https://doi.org/10.1016/j.jep.2022.115410>.
- [45] Zheng Y, Zhou X, Wang C, Zhang J, Chang D, Zhuang S, et al. Effect of dendrobium mixture in alleviating diabetic cognitive impairment associated with regulating gut microbiota. *Biomed Pharmacother* 2022;149:112891. doi: <https://doi.org/10.1016/j.biopha.2022.112891>.
- [46] Wang D, Wang L, Han L, Wang B, Shi R, Ye J, et al. Leucine-restricted diet ameliorates obesity-linked cognitive deficits: involvement of the microbiota-gut-brain axis. *J Agric Food Chem* 2023;71(24):9404–18. doi: <https://doi.org/10.1021/acs.jafc.3c01524>.
- [47] Zhuang ZQ, Shen LL, Li WW, Fu X, Zeng F, Gui L, et al. Gut microbiota is altered in patients with Alzheimer's disease. *J Alzheimers Dis* 2018;63(4):1337–46. doi: <https://doi.org/10.3233/jad-180176>.
- [48] Chen L, Xu X, Wu X, Cao H, Li X, Hou Z, et al. A comparison of the composition and functions of the oral and gut microbiotas in Alzheimer's patients. *Front Cell Infect Microbiol* 2022;12:942460. doi: <https://doi.org/10.3389/fcimb.2022.942460>.
- [49] Lu S, Yang Y, Xu Q, Wang S, Yu J, Zhang B, et al. Gut microbiota and targeted biomarkers analysis in patients with cognitive impairment. *Front Neurol* 2022;13:84403. doi: <https://doi.org/10.3389/fneur.2022.834403>.
- [50] Upadhyay P, Tyagi A, Agrawal S, Kumar A, Gupta S. Bidirectional effect of Triphala on modulating gut-brain axis to improve cognition in the murine model of Alzheimer's disease. *Mol Nutr Food Res* 2023:e2300104. doi: <https://doi.org/10.1002/mnfr.202300104>.
- [51] Reemst K, Tims S, Yam KY, Mischke M, Knol J, Brul S, et al. The role of the gut microbiota in the effects of early-life stress and dietary fatty acids on later-life central and metabolic outcomes in mice. *mSystems* 2022;7(3):e0018022. DOI: 10.1128/mSystems.00180-22.
- [52] Khedr EM, Omeran N, Karam-Allah Ramadan H, Ahmed GK, Abdelwarith AM. Alteration of gut microbiota in Alzheimer's disease and their relation to the cognitive impairment. *J Alzheimers Dis* 2022;88(3):1103–14. doi: <https://doi.org/10.3233/jad-220176>.
- [53] Kameno K, Hasegawa Y, Hayashi K, Takemoto Y, Uchikawa H, Mukasa A, et al. Loss of body weight in old 5xFAD mice and the alteration of gut microbiota composition. *Exp Gerontol* 2022;166:111885. doi: <https://doi.org/10.1016/j.exger.2022.111885>.
- [54] Ma J, Song XQ. Correlation between cognitive impairment and metabolic imbalance of gut microbiota in patients with schizophrenia. *World J Psychiatry* 2023;13(10):724–31. doi: <https://doi.org/10.5498/wjp.v13.i10.724>.
- [55] Zhang P, Wang T, Zhu X, Feng L, Wang J, Li Y, et al. Jiedu Yizhi formula improves cognitive function by regulating the gut dysbiosis and TLR4/NF- κ B signaling pathway. *Neuropsychiatr Dis Treat* 2023;19:49–62. doi: <https://doi.org/10.2147/ndt.S393773>.
- [56] Hua D, Li S, Li S, Wang X, Wang Y, Xie Z, et al. Gut microbiome and plasma metabolome signatures in middle-aged mice with cognitive dysfunction induced by chronic neuropathic pain. *Front Mol Neurosci* 2021;14:806700. doi: <https://doi.org/10.3389/fnmol.2021.806700>.
- [57] Cilia R, Piatti M, Cereda E, Bolliri C, Caronni S, Ferri V, et al. Does gut microbiota influence the course of Parkinson's disease? A 3-year prospective exploratory study in de novo patients. *J Parkinsons Dis* 2021;11(1):159–70. doi: <https://doi.org/10.3233/jpd-202297>.
- [58] Su A, Yang W, Zhao L, Pei F, Yuan B, Zhong L, et al. Flammulina velutipes polysaccharides improve scopolamine-induced learning and memory impairment in mice by modulating gut microbiota composition. *Food Funct* 2018;9(3):1424–32. doi: <https://doi.org/10.1039/c7fo01991b>.
- [59] Putri SSF, Irfannuddin I, Murti K, Kesuma Y, Darmawan H, Koibuchi N. The role of gut microbiota on cognitive development in rodents: A meta-analysis. *J Physiol Sci* 2023;73(1):10. doi: <https://doi.org/10.1186/s12576-023-00869-1>.
- [60] McLeod A, Penálver Bernabe B, Xia Y, Sanchez-Flack J, Lamar M, Schiffer L, et al. Comparing the gut microbiome of obese, African American, older adults with and without mild cognitive impairment. *PLoS One* 2023;18(2):e0280211.
- [61] Gao Q, Li D, Wang Y, Zhao C, Li M, Xiao J, et al. Analysis of intestinal flora and cognitive function in maintenance hemodialysis patients using combined 16S ribosome DNA and shotgun metagenome sequencing. *Aging Clin Exp Res* 2024;36(1):28. doi: <https://doi.org/10.1007/s40520-023-02645-y>.
- [62] Bloemendaal M, Szopinska-Tokov J, Belzer C, Boverhoff D, Papalini S, Michels F, et al. Probiotics-induced changes in gut microbial composition and its effects on cognitive performance after stress: exploratory analyses. *Transl Psychiatry* 2021;11(1):300. doi: <https://doi.org/10.1038/s41398-021-01404-9>.
- [63] Yang P, Huang S, Luo Z, Zhou S, Zhang C, Zhu Y, et al. Radix Bupleuri aqueous extract attenuates MK801-induced schizophrenia-like symptoms in mice: Participation of intestinal flora. *Biomed Pharmacother* 2024;172:116267. doi: <https://doi.org/10.1016/j.biopha.2024.116267>.
- [64] Luo XX, Yang C, Zhan GF, Li S, Hua DY, Luo AL, et al. Whole brain radiotherapy induces cognitive dysfunction in mice: key role of gut microbiota. *Psychopharmacology* 2020;237(7):2089–101. doi: <https://doi.org/10.1007/s00213-020-05520-0>.
- [65] Hu X, Mao Y, Luo F, Wang X. Association between post-stroke cognitive impairment and gut microbiota: A PRISMA-compliant systematic review and meta-analysis. *Medicine (Baltimore)* 2023;102(35):e34764.
- [66] Salazar N, Arbolea S, Fernández-Navarro T, de Los Reyes-Gavilán CG, González S, Gueimonde M. Age-associated changes in gut microbiota and dietary components related with the immune system in adulthood and old age: A cross-sectional study. *Nutrients* 2019;11(8):1765. doi: <https://doi.org/10.3390/nu11081765>.
- [67] Zhang Y, Sun D, Zhao X, Luo Y, Yu H, Zhou Y, et al. Bacteroides fragilis prevents aging-related atrial fibrillation in rats via regulatory T cells-mediated regulation of inflammation. *Pharmacol Res* 2022;177:106141. doi: <https://doi.org/10.1016/j.phrs.2022.106141>.
- [68] Aljumaah MR, Bhatia U, Roach J, Gunstad J, Azcarate Peril MA. The gut microbiome, mild cognitive impairment, and probiotics: A randomized clinical trial in middle-aged and older adults. *Clin Nutr* 2022;41(11):2565–76. doi: <https://doi.org/10.1016/j.clnu.2022.09.012>.
- [69] Pan Q, Li YQ, Guo K, Xue M, Gan Y, Wang K, et al. Elderly patients with mild cognitive impairment exhibit altered gut microbiota profiles. *J Immunol Res* 2021;2021:5578958. doi: <https://doi.org/10.1155/2021/5578958>.
- [70] Tamana SK, Tun HM, Konya T, Chari RS, Field CJ, Guttman DS, et al. Bacteroides-dominant gut microbiome of late infancy is associated with enhanced neurodevelopment. *Gut Microbes* 2021;13(1):1–17. doi: <https://doi.org/10.1080/19490976.2021.1930875>.
- [71] Park SH, Lee JH, Kim JS, Kim TJ, Shin J, Im JH, et al. Fecal microbiota transplantation can improve cognition in patients with cognitive decline and Clostridioides difficile infection. *Aging (Albany NY)* 2022;14(16):6449–66. doi: <https://doi.org/10.18632/aging.204230>.
- [72] Wang Q, Song YX, Wu XD, Luo YG, Miao R, Yu XM, et al. Gut microbiota and cognitive performance: A bidirectional two-sample Mendelian randomization. *J Affect Disord* 2024:35338–47. doi: <https://doi.org/10.1016/j.jad.2024.02.083>.
- [73] Chen C, Liao J, Xia Y, Liu X, Jones R, Haran J, et al. Gut microbiota regulate Alzheimer's disease pathologies and cognitive disorders via PUFA-associated neuroinflammation. *Gut* 2022;71(11):2233–52. doi: <https://doi.org/10.1136/gutjnl-2021-326269>.
- [74] Saji N, Murotani K, Hisada T, Tsuduki T, Sugimoto T, Kimura A, et al. The relationship between the gut microbiome and mild cognitive impairment in patients without dementia: A cross-sectional study conducted in Japan. *Sci Rep* 2019;9(1):19227. doi: <https://doi.org/10.1038/s41598-019-55851-y>.
- [75] Chen Y, Ouyang Y, Chen X, Chen R, Ruan Q, Farag MA, et al. Hypoglycaemic and anti-ageing activities of green alga Ulva lactuca polysaccharide via gut microbiota in ageing-associated diabetic mice. *Int J Biol Macromol* 2022;212:97. doi: <https://doi.org/10.1016/j.ijbiomac.2022.05.109>.
- [76] Qian X, Hai W, Chen S, Zhang M, Jiang X, Tang H. Multi-omics data reveals aberrant gut microbiota-host glycerophospholipid metabolism in association with neuroinflammation in APP/PS1 mice. *Gut Microbes* 2023;15(2):2282790. doi: <https://doi.org/10.1080/19490976.2023.2282790>.
- [77] Xu Y, Yang Y, Li B, Xie Y, Shi Y, Le G. Dietary methionine restriction improves gut microbiota composition and prevents cognitive impairment in D-galactose-induced aging mice. *Food Funct* 2022;13(24):12896–914. doi: <https://doi.org/10.1039/d2fo03366f>.
- [78] Akbari E, Asemi Z, Daneshvar Kakhaki R, Bahmani F, Kouchaki E, Tamtaji OR, et al. Effect of probiotic supplementation on cognitive function and metabolic status in Alzheimer's disease: A randomized, double-blind and controlled trial. *Front Aging Neurosci* 2016;8:256. doi: <https://doi.org/10.3389/fnagi.2016.00256>.
- [79] Xiao Y, Yang C, Yu L, Tian F, Wu Y, Zhao J, et al. Human gut-derived B. longum subsp. longum strains protect against aging in a D-galactose-induced aging mouse model. *Microbiome* 2021;9(1):180. doi: <https://doi.org/10.1186/s40168-021-01108-8>.
- [80] Shi S, Zhang Q, Sang Y, Ge S, Wang Q, Wang R, et al. Probiotic bifidobacterium longum bb68s improves cognitive functions in healthy older adults: a randomized, double-blind, placebo-controlled trial. *Nutrients* 2022;15(1):51. doi: <https://doi.org/10.3390/nu15010051>.
- [81] Robertson RC, Seira Oriach C, Murphy K, Moloney GM, Cryan JF, Dinan TG, et al. Omega-3 polyunsaturated fatty acids critically regulate behaviour and gut microbiota development in adolescence and adulthood. *Brain Behav Immun* 2017;59:21–37. doi: <https://doi.org/10.1016/j.bbi.2016.07.145>.
- [82] Azuma N, Mawatari T, Saito Y, Tsukamoto M, Sampei M, Iwama Y. Effect of continuous ingestion of bifidobacteria and dietary fiber on improvement in cognitive function: A randomized, double-blind, placebo-controlled trial. *Nutrients* 2023;15(19):4175. doi: <https://doi.org/10.3390/nu15194175>.
- [83] Ni Y, Yang X, Zheng L, Wang Z, Wu L, Jiang J, et al. Lactobacillus and bifidobacterium improves physiological function and cognitive ability in aged

- mice by the regulation of gut microbiota. *Mol Nutr Food Res* 2019;63(22): e1900603.
- [84] O'Hagan C, Li JV, Marchesi JR, Plummer S, Garaiova I, Good MA. Long-term multi-species Lactobacillus and Bifidobacterium dietary supplement enhances memory and changes regional brain metabolites in middle-aged rats. *Neurobiol Learn Mem* 2017;14436–47. doi: <https://doi.org/10.1016/j.nlm.2017.05.015>.
- [85] Abdelhamid M, Zhou C, Jung CG, Michikawa M. Probiotic Bifidobacterium breve MCC1274 mitigates Alzheimer's disease-related pathologies in wild-type mice. *Nutrients* 2022;14(12):2543. doi: <https://doi.org/10.3390/nu14122543>.
- [86] Zhu G, Zhao J, Zhang H, Chen W, Wang G. Administration of Bifidobacterium breve improves the brain function of A β (1–42)-treated mice via the modulation of the gut microbiome. *Nutrients* 2021;13(5):1602. doi: <https://doi.org/10.3390/nu13051602>.
- [87] Araki A, Yoshimura Y, Sakurai T, Umegaki H, Kamada C, Iimuro S, et al. Low intakes of carotene, vitamin B(2), pantothenate and calcium predict cognitive decline among elderly patients with diabetes mellitus: The Japanese elderly diabetes intervention trial. *Geriatr Gerontol Int* 2017;17(8):1168–75. doi: <https://doi.org/10.1111/ggi.12843>.
- [88] Liu FC, Cheng ML, Lo CJ, Hsu WC, Lin G, Lin HT. Exploring the aging process of cognitively healthy adults by analyzing cerebrospinal fluid metabolomics using liquid chromatography-tandem mass spectrometry. *BMC Geriatr* 2023;23(1):217. doi: <https://doi.org/10.1186/s12877-023-03939-6>.
- [89] Kodam P, Sai Swaroop R, Pradhan SS, Sivaramakrishnan V, Vadrevu R. Integrated multi-omics analysis of Alzheimer's disease shows molecular signatures associated with disease progression and potential therapeutic targets. *Sci Rep* 2023;13(1):3695. doi: <https://doi.org/10.1038/s41598-023-30892-6>.
- [90] Moreno DJM, M. Cognitive improvement in mild to moderate Alzheimer's dementia after treatment with the acetylcholine precursor choline alfoscerate: a multicenter, double-blind, randomized, placebo-controlled trial. *Clin Ther* 2003;25(1):178–93. doi: [https://doi.org/10.1016/s0149-2918\(03\)90023-3](https://doi.org/10.1016/s0149-2918(03)90023-3).
- [91] Narukawa M, Kamiyoshihara A, Izu H, Fujii T, Matsubara K, Misaka T. Efficacy of long-term feeding of α -glycerophosphocholine for aging-related phenomena in old mice. *Gerontology* 2020;66(3):275–85. doi: <https://doi.org/10.1159/000504962>.
- [92] Sagaro GG, Traini E, Amenta F. Activity of choline alfoscerate on adult-onset cognitive dysfunctions: A systematic review and meta-analysis. *J Alzheimers Dis* 2023;92(1):59–70. doi: <https://doi.org/10.3233/jad-221189>.
- [93] Guan W, Xu DW, Ji CH, Wang CN, Liu Y, Tang WQ, et al. Hippocampal miR-206-3p participates in the pathogenesis of depression via regulating the expression of BDNF. *Pharmacol Res* 2021;174105932. doi: <https://doi.org/10.1016/j.phrs.2021.105932>.
- [94] Peng D, Wang Y, Xiao Y, Peng M, Mai W, Hu B, et al. Extracellular vesicles derived from astrocyte-treated with hA β GF(14–154) attenuate Alzheimer phenotype in AD mice. *Theranostics* 2022;12(8):3862–81. doi: <https://doi.org/10.7150/thno.70951>.
- [95] Kondo M, Yamada H, Munetsuna E, Yamazaki M, Hatta T, Iwahara A, et al. Associations of serum microRNA-20a, -27a, and -103a with cognitive function in a Japanese population: The Yakumo study. *Arch Gerontol Geriatr* 2019;82155–60. doi: <https://doi.org/10.1016/j.archger.2019.01.007>.
- [96] Hashemi KS, Aliabadi MK, Mehrara A, Talebi E, Hemmati AA, Rezaeiyari RD, et al. A meta-analysis of microarray datasets to identify biological regulatory networks in Alzheimer's disease. *Front Genet* 2023;141225196. doi: <https://doi.org/10.3389/fgene.2023.1225196>.
- [97] Da Silva FC, Rode MP, Vietta GG, Iop RDR, Creczynski-Pasa TB, Martin AS, et al. Expression levels of specific microRNAs are increased after exercise and are associated with cognitive improvement in Parkinson's disease. *Mol Med Rep* 2021;24(2). doi: <https://doi.org/10.3892/mmr.2021.12257>.
- [98] Pereira JB, Janelidze S, Strandberg O, Whelan CD, Zetterberg H, Blennow K, et al. Microglial activation protects against accumulation of tau aggregates in nondemented individuals with underlying Alzheimer's disease pathology. *Nat Aging* 2022;2(12):1138–44. doi: <https://doi.org/10.1038/s43587-022-00310-z>.
- [99] Su Q, Ren YH, Liu GW, Gao YP, Zhang JX, Zhang JN, et al. Trichostatin A relieves anxiety-and depression-like symptoms in APP/PS1 mice. *Front Pharmacol* 2024;151333235. doi: <https://doi.org/10.3389/fphar.2024.1333235>.
- [100] Yoon H, Radulovic M, Wu J, Blaber SI, Blaber M, Fehlings MG, et al. Kallikrein 6 signals through PAR1 and PAR2 to promote neuron injury and exacerbate glutamate neurotoxicity. *J Neurochem* 2013;127(2):283–98. doi: <https://doi.org/10.1111/jnc.12293>.
- [101] Patra K, Soosaipillai A, Sando SB, Lauridsen C, Berge G, Møller I, et al. Assessment of kallikrein 6 as a cross-sectional and longitudinal biomarker for Alzheimer's disease. *Alzheimers Res Ther* 2018;10(1):9. doi: <https://doi.org/10.1186/s13195-018-0336-4>.
- [102] Zhang Z, Jhaveri D, Sharmin S, Harvey TJ, Dawson PA, Piper M, et al. Cell-extrinsic requirement for sulfate in regulating hippocampal neurogenesis. *Biol Open* 2020;9(7). doi: <https://doi.org/10.1242/bio.053132>.
- [103] Wang Y, Lin Y, Wang L, Zhan H, Luo X, Zeng Y, et al. TREM2 ameliorates neuroinflammatory response and cognitive impairment via PI3K/AKT/FoxO3a signaling pathway in Alzheimer's disease mice. *Aging (Albany NY)* 2020;12(20):20862–79. doi: <https://doi.org/10.18632/aging.104104>.
- [104] Fassler M, Rappaport MS, Cuño CB, George J. Engagement of TREM2 by a novel monoclonal antibody induces activation of microglia and improves cognitive function in Alzheimer's disease models. *J Neuroinflammation* 2021;18(1):19. doi: <https://doi.org/10.1186/s12974-020-01980-5>.
- [105] Frey S, Eichler A, Stonawski V, Kriebel J, Wahl S, Gallati S, et al. Prenatal alcohol exposure is associated with adverse cognitive effects and distinct whole-genome dna methylation patterns in primary school children. *Front Behav Neurosci* 2018;12125. doi: <https://doi.org/10.3389/fnbeh.2018.00125>.
- [106] Liang T, Ju H, Zhou Y, Yang Y, Shi Y, Fang H. Inhibition of glycogen synthase kinase 3 β improves cognitive function in aged mice by upregulating claudin presences in cerebral endothelial cells. *Acta Biochim Biophys Sin (Shanghai)* 2020;52(4):363–70. doi: <https://doi.org/10.1093/abbs/gmaa002>.
- [107] Chen Y, Zheng Z, Mei A, Huang H, Lin F. Claudin-1 and Claudin-3 as molecular regulators of myelination in leukoaraiosis patients. *Clinics (Sao Paulo)* 2021;76e2167. doi: <https://doi.org/10.6061/clinics/2021/e2167>.
- [108] Liu S, Ando F, Fujita Y, Liu J, Maeda T, Shen X, et al. A clinical dose of angiotensin-converting enzyme (ACE) inhibitor and heterozygous ACE deletion exacerbate Alzheimer's disease pathology in mice. *J Biol Chem* 2019;294(25):9760–70. doi: <https://doi.org/10.1074/jbc.RA118.006420>.
- [109] Hsieh SW, Liu MW, Huang LC, Wu MN, Yang YH. The impact of angiotensin-converting enzyme gene on behavioral and psychological symptoms of dementia in Alzheimer's disease. *Curr Alzheimer Res* 2019;16(14):1269–75. doi: <https://doi.org/10.2174/1567205017666200103114550>.
- [110] Yip AG, Brayne C, Easton D, Rubinstein DC. An investigation of ACE as a risk factor for dementia and cognitive decline in the general population. *J Med Genet* 2002;39(6):403–6. doi: <https://doi.org/10.1136/img.39.6.403>.
- [111] Deng Z, Jiang J, Wang J, Pan D, Zhu Y, Li H, et al. Angiotensin receptor blockers are associated with a lower risk of progression from mild cognitive impairment to dementia. *Hypertension* 2022;79(10):2159–69. doi: <https://doi.org/10.1161/hypertensionaha.122.19378>.
- [112] Loerch PM, Lu T, Dakin KA, Vann JM, Isaacs A, Geula C, et al. Evolution of the aging brain transcriptome and synaptic regulation. *PLoS One* 2008;3(10): e3329.
- [113] Rehman H, Ang TFA, Tao Q, Espenilla AL, Au R, Farrer LA, et al. Comparison of commonly measured plasma and cerebrospinal fluid proteins and their significance for the characterization of cognitive impairment status. *J Alzheimers Dis* 2024;97(2):621–33. doi: <https://doi.org/10.3233/jad-230837>.
- [114] Zou S, Zhang J, Chen W. Subtypes based on six apolipoproteins in non-demented elderly are associated with cognitive decline and subsequent tau accumulation in cerebrospinal fluid. *J Alzheimers Dis* 2019;72(2):413–23. doi: <https://doi.org/10.3233/jad-190314>.
- [115] Carvalho K, Scharzt ND, Balderrama-Gutierrez G, Liang HY, Chu SH, Selvan P, et al. Modulation of C5a–C5aR1 signaling alters the dynamics of AD progression. *J Neuroinflammation* 2022;19(1):178. doi: <https://doi.org/10.1186/s12974-022-02539-2>.
- [116] Liang Y, Kang X, Zhang H, Xu H, Wu X. Knockdown and inhibition of hippocampal GPR17 attenuates lipopolysaccharide-induced cognitive impairment in mice. *J Neuroinflammation* 2023;20(1):271. doi: <https://doi.org/10.1186/s12974-023-02958-9>.
- [117] Wang Y, Jiang A, Yan J, Wen D, Gu N, Li Z, et al. Inhibition of GPR17/ID2 axis improve remyelination and cognitive recovery after SAH by mediating OPC differentiation in rat model. *Transl Stroke Res* 2023. doi: <https://doi.org/10.1007/s12975-023-01201-0>.
- [118] Pettit NL, Yap EL, Greenberg ME, Harvey CD. Fos ensembles encode and shape stable spatial maps in the hippocampus. *Nature* 2022;609(7926):327–34. doi: <https://doi.org/10.1038/s41586-022-05113-1>.
- [119] Lanke V, Moolamalla STR, Roy D, Vinod PK. Integrative analysis of hippocampus gene expression profiles identifies network alterations in aging and Alzheimer's Disease. *Front Aging Neurosci* 2018;10153. doi: <https://doi.org/10.3389/fnagi.2018.00153>.
- [120] Weber M, Wu T, Hanson JE, Alam NM, Solano H, Ngu H, et al. Cognitive deficits, changes in synaptic function, and brain pathology in a mouse model of normal aging(1,2,3). *eNeuro* 2015;2(5):ENEURO.0047-15. DOI: 10.1523/eneuro.0047-15.2015.
- [121] Shevtsova O, Tan YF, Merkley CM, Winocur G, Wojtowicz JM. Early-age running enhances activity of adult-born dentate granule neurons following learning in rats. *eNeuro* 2017;4(4):ENEURO.0237-17.2017. doi: <https://doi.org/10.1523/eneuro.0237-17.2017>.
- [122] Trettel CDS, Pelozin BRA, Barros MP, Bachi ALL, Braga PGS, Momesso CM, et al. Irisin: An anti-inflammatory exerkine in aging and redox-mediated comorbidities. *Front Endocrinol (Lausanne)* 2023;141106529. doi: <https://doi.org/10.3389/fendo.2023.1106529>.
- [123] Zhu W, Sahar NE, Javaid HMA, Pak ES, Liang G, Wang Y, et al. Exercise-induced irisin decreases inflammation and improves NAFLD by competitive binding with MD2. *Cells* 2021;10(12):3306. doi: <https://doi.org/10.3390/cells10123306>.
- [124] Cribbs DH, Berchtold NC, Perreau V, Coleman PD, Rogers J, Tenner AJ, et al. Extensive innate immune gene activation accompanies brain aging, increasing vulnerability to cognitive decline and neurodegeneration: a microarray study. *J Neuroinflammation* 2012;9179. doi: <https://doi.org/10.1186/1742-2094-9-179>.
- [125] Calvo-Rodríguez M, de la Fuente C, García-Durillo M, García-Rodríguez C, Villalobos C, Núñez L. Aging and amyloid β oligomers enhance TLR4 expression, LPS-induced Ca(2+) responses, and neuron cell death in cultured rat hippocampal neurons. *J Neuroinflammation* 2017;14(1):24. doi: <https://doi.org/10.1186/s12974-017-0802-0>.

- [126] Potter OV, Giedraitis ME, Johnson CD, Cox MN, Kohman RA. Young and aged TLR4 deficient mice show sex-dependent enhancements in spatial memory and alterations in interleukin-1 related genes. *Brain Behav Immun* 2019;7637–47. doi: <https://doi.org/10.1016/j.bbi.2018.10.010>.
- [127] Connolly MG, Yost OL, Potter OV, Giedraitis ME, Kohman RA. Toll-like receptor 4 differentially regulates adult hippocampal neurogenesis in an age- and sex-dependent manner. *Hippocampus* 2020;30(9):958–69. doi: <https://doi.org/10.1002/hipo.23209>.
- [128] Denver P, English A, McClean PL. Inflammation, insulin signaling and cognitive function in aged APP/PS1 mice. *Brain Behav Immun* 2018;70423–34. doi: <https://doi.org/10.1016/j.bbi.2018.03.032>.
- [129] Moreno-Cugnon L, Revuelta M, Arrizabalaga O, Colie S, Moreno-Valladares M, Jimenez-Blasco D, et al. Neuronal p38 α mediates age-associated neural stem cell exhaustion and cognitive decline. *Aging Cell* 2019;18(6):e13044.
- [130] Oskouei Z, Mehri S, Kalalinia F, Hosseinzadeh H. Evaluation of the effect of thymoquinone in d-galactose-induced memory impairments in rats: Role of MAPK, oxidative stress, and neuroinflammation pathways and telomere length. *Phytother Res* 2021;35(4):2252–66. doi: <https://doi.org/10.1002/ptr.6982>.
- [131] Cardoso FDS, França EF, Serra FT, Victorino AB, de Almeida AA, Fernandes J, et al. Aerobic exercise reduces hippocampal ERK and p38 activation and improves memory of middle-aged rats. *Hippocampus* 2017;27(8):899–905. doi: <https://doi.org/10.1002/hipo.22740>.
- [132] Kim KY, Kwak S, Ha J, Oh DJ, Kim M, Cho SY, et al. Loss of association between plasma irisin levels and cognition in Alzheimer's disease. *Psychoneuroendocrinology* 2022;136105624. doi: <https://doi.org/10.1016/j.psyneuen.2021.105624>.
- [133] Wu H, Zhang Q, Xu P, Chen J, Duan L, Xu F, et al. Nattokinase promotes post-stroke neurogenesis and cognition recovery via increasing circulating irisin. *J Agric Food Chem* 2023;71(30):11418–28. doi: <https://doi.org/10.1021/acs.jafc.2c08718>.
- [134] Song D, Chen X, Zhou N, Yuan Y, Geng S, Zhang C, et al. Low-intensity pulsed ultrasound triggers a beneficial neuromodulation in dementia mice with chronic cerebral hypoperfusion via activation of hippocampal Fndc5/irisin signaling. *J Transl Med* 2023;21(1):139. doi: <https://doi.org/10.1186/s12967-022-03824-7>.
- [135] Lee H, Kim SY, Lim Y. Solanum melongena extract supplementation protected skeletal muscle and brain damage by regulation of BDNF/PGC1 α /irisin pathway via brain function-related myokines in high-fat diet induced obese mice. *J Nutr Biochem* 2024;124109537. doi: <https://doi.org/10.1016/j.jnutbio.2023.109537>.